



Reports of Suspected Adverse Reaction to COVID-19 Vaccines (01 March to 07 November 2021)

Contents

About the report	2
Summary	3
COVID-19 vaccines with Emergency Use Authorization in the Philippines	4
Statistics regarding reports of suspected adverse reactions	4
Distribution of reports of adverse reaction for each vaccine.....	5
Demographics	5
Vaccination in adolescent population	6
Reports of suspected serious adverse reaction	7
Number of suspected adverse reactions per category	11
Reactions to inactivated vaccines	12
Reactions to non-replicating viral vector vaccines	14
Reactions to mRNA vaccines	17
Outcome of suspected adverse reaction	19
Reporting of suspected adverse reactions following vaccination	19



About the report

- A summary is presented below of all received suspected adverse reaction reports following COVID-19 vaccination from 01 March 2021, the date when the first vaccine became available, up to 07 November 2021.
- Seven (7) vaccines under Emergency Use Authorization (EUA) are currently being used in the vaccination program: CoronaVac, COVID-19 Vaccine AstraZeneca, Sputnik V, Comirnaty, COVID-19 Vaccine Moderna, Janssen COVID-19 Vaccine, and COVID-19 Vaccine Sinopharm.
- Data are based on VigiFlow, the national database of adverse reactions in the Philippines. It includes reports from various epidemiology surveillance units (ESUs) of the Department of Health (DOH), hospitals, patients/consumers, and EUA holders.
- Symptoms or diseases that occur after vaccination are reported if there is a *suspicion* of a possible link. However, it cannot be assumed that there is a causal relationship between the suspected adverse reaction and the vaccine.
- This report contains all suspected adverse reactions regardless of any possible causal relationship.
- Additional information may become available in individual case reports at any time, which may change the assessment and figures presented.
- Adverse reaction reports are necessary for the safety assessment of the vaccines, making sure that the benefits always outweigh the risks.
- Reports are constantly reviewed and monitored for the possible emergence/identification of unknown adverse reactions also known as signal. If a signal is identified, investigations, regulatory actions, and timely communication is performed by the FDA.
- A weekly report is published to summarize reported adverse reactions to the COVID-19 vaccines.

Summary

This report is based on an assessment of adverse reaction reports received by 07 November 2021. As per benefit-risk assessment, these reports do not provide a basis for revising the current recommendations regarding the use of COVID-19 vaccines.

A report of adverse reaction does not necessarily mean that the vaccine caused the reactions. A mere suspicion may also be reported. Undiagnosed illness, underlying comorbidities, and pre-existing medical conditions unrelated to vaccination can be factors in reporting adverse reactions. The relative numbers should not be used to compare the safety of different vaccines.

Like any other vaccines, COVID-19 vaccines may cause adverse reactions in some people. It is possible for several persons to experience the same adverse event but for the report to be serious for one person and non-serious for another person. Most of the reported reactions are generally in line with what is described in the product information and labels. Such reports are minor adverse reactions, which include body pain, chills, fatigue, fever, headache, nausea, and pain in the injection site. These usually appear on the first or second day of vaccination and may last for 2-3 days. Most people tolerate these adverse reactions while others experience greater discomfort.

Serious adverse reactions have also been reported. The FDA together with other public health partners are continuously monitoring the adverse experience as more people are being vaccinated with COVID-19 vaccines. Such monitoring will provide reassurance that the vaccines are safe and effective for use.

Considering the post-authorization experience on the use of COVID-19 Vaccine AstraZeneca and Janssen COVID-19 Vaccine of other countries, information on the very rare and serious adverse events of thrombosis and thrombocytopenia and in some cases accompanied by bleeding have been revised under the special warning and precautions for use in their product labels. In addition, capillary leak syndrome and Guillain-Barré syndrome were added as a rare new side effect.

The label of mRNA vaccines Moderna and Comirnaty (Pfizer-BioNTech) have been revised to include imposition of the European Medicines Agency and the USFDA to include safety information on myocarditis and pericarditis.

The immunization program expanded its coverage to include adolescent individuals (12 to 17 years old). Pfizer and Moderna are the only vaccines with emergency use authorization for the said population. The roll-out started on the second week of October. Likewise, monitoring of suspected adverse reactions has extended to include such population.

Seven (7) vaccines are currently used in the immunization program. These include CoronaVac, COVID-19 Vaccine AstraZeneca, Sputnik V, Comirnaty, COVID-19 Vaccine Moderna, Janssen COVID-19 Vaccine, and COVID-19 Vaccine Sinopharm. Supplies of vaccines are either procured by the government and/or private sector or supplied under the COVAX facility.

COVID-19 vaccines with Emergency Use Authorization in the Philippines

At present, the FDA granted [eight \(8\) COVID-19 vaccines with emergency use authorization](#):

- Pfizer-BioNTech COVID-19 Vaccine (Comirnaty)
- ChAdOx1-S [recombinant] (COVID-19 Vaccine AstraZeneca)
- SARS-CoV-2 Vaccine (Vero Cell), Inactivated (CoronaVac)
- Gam-COVID-Vac (Sputnik V & Sputnik Light)
- Ad26.COV2-S [recombinant] (Janssen COVID-19 Vaccine)
- Whole Virion Inactivated Corona Virus (Covaxin)
- COVID-19 mRNA Vaccine [nucleoside modified] (COVID-19 Vaccine Moderna)
- Inactivated COVID-19 Vaccine (Vero Cell) (COVID-19 Vaccine Sinopharm BIBP/Wuhan)

Various vaccine platforms have been approved for use in the Philippines. Comirnaty and COVID-19 Vaccine Moderna are mRNA vaccines; COVID-19 Vaccine AstraZeneca and Janssen COVID-19 Vaccine are non-replicating viral vector vaccines while Sputnik V uses the same technology having two (2) different (dose) components of viral vectors; and CoronaVac, Covaxin, and COVID-19 Vaccine Sinopharm BIBP/Wuhan are inactivated vaccines. All are administered in two doses within an interval of a few weeks except for Janssen COVID-19 Vaccine and Sputnik Light, which is administered as a single-dose.

Statistics regarding reports of suspected adverse reactions

As of 07 November 2021, more than 64.1 million doses of COVID-19 vaccines (either CoronaVac, COVID-19 Vaccine AstraZeneca, Sputnik V, Comirnaty, COVID-19 Vaccine Moderna, Janssen COVID-19 Vaccine, or COVID-19 Vaccine Sinopharm) were already administered. Over 29.4 million individuals are now fully vaccinated (either given a single-dose or 2-dose vaccine course) while more than 8.8 million are partly vaccinated waiting for their second dose to be administered. A total of 73,281 suspected adverse reaction reports were received, evaluated, and analyzed by the FDA. To disaggregate, 28,899 have been reported for CoronaVac, 32,454 for COVID-19 Vaccine AstraZeneca, 743 for Sputnik V, 5,256 for Comirnaty, 2,343 for COVID-19 Vaccine Moderna, 3,477 for Janssen COVID-19 Vaccine, and 109 for COVID-19 Vaccine Sinopharm.

Table 1. Data on vaccination and suspected adverse reaction reports.

Indicators	Value
Total number of doses administered	64,195,936
No. of fully vaccinated individuals	29,477,961
No. of individuals partly vaccinated	8,862,889
No. of suspected adverse reaction reports	73,281 (0.11% of doses administered)
No. of suspected serious adverse reaction reports	3,601 (0.006% of doses administered)

Distribution of reports of adverse reactions for each vaccine

Data shown below are cumulative reports from the start of the vaccination program on 01 March 2021 up until 07 November 2021.

Table 2. Distribution of reports of adverse reactions for each vaccine

Vaccine	Date started	Total vaccine doses administered ^b	Number of fully vaccinated individuals ^b	Number of individuals partly vaccinated	Total number of reports ^a	Reports of non-serious events	Reports of serious events
CoronaVac	01 Mar 2021	32,063,369	14,741,907	2,579,555	28,899	27,184	1,715
AstraZeneca	07 Mar 2021	9,205,925	3,959,183	1,287,559	32,454	31,472	982
Sputnik V	04 May 2021	564,585	251,013	62,559	743	719	24
Comirnaty	13 May 2021	11,933,502	4,396,861	3,139,780	5,256	4,960	296
Moderna	30 June 2021	5,956,059	2,118,716	1,718,627	2,343	2,203	140
Janssen	20 July 2021	3,622,875	3,622,875	-	3,477	3,041	436
Sinopharm	25 Aug 2021	849,621	387,406	74,809	109	101	8
TOTAL		64,195,936	29,477,961	8,862,889	73,281	69,680	3,601

Data source: ^aVigiFlow, ^bNVOC daily report as 6PM, 07 November 2021

Notes: Additional information may become available in individual cases, which may change the figures presented

^aAn individual is considered partly vaccinated if they have received only one dose of a two-dose vaccine course. An individual is considered fully vaccinated if they have received a single-dose vaccine or both doses of a two-dose vaccine

^cData concerning various vaccines are not directly comparable. COVID-19 vaccines profile varies, they have not been used for equal periods of time and they have been administered to number of people with different profiles including various age and sex.

Demographics

The figures below provide a descriptive overview of the population who have experienced adverse reactions to COVID-19 vaccines. Figure 1 and Figure 2 shows the distribution of reports by gender and age. Click [here](#) to show disaggregated data.




Figure 1. Report distribution by gender




Figure 2. Report distribution by age

The early phase of the vaccination program is intended for the frontline health workers, thus, the high number in the female and younger population might be associated with the data that our health system is dominated by female (75%) and young adults under the age of 35 (65%).¹ An increasing number of reports from the age group 40 years and above have been observed in the past few weeks of the vaccination program. This may be attributed to the coverage of priority groups of senior citizens and individuals with comorbidities.

Relative to the inclusion of the frontline personnel in the priority groups, the observed increasing number of reports in the male population may be attributed to the vaccine coverage and statistics that more males are employed than females (6 in every 10).²

¹ Human Resource for Health in the Time of the COVID-19 Pandemic: Does the Philippines Have Enough? <https://www.drdf.org.ph/sites/default/files/pdf/COVID-19-Research-Brief-08.pdf>

² Employment situation in July 2018, Philippine Statistics Authority <https://psa.gov.ph/statistics/survey/labor-and-employment/labor-force-survey/title/Employment%20Situation%20in%20July%202018>

Vaccination in adolescent population

The roll out for vaccinating adolescent population started last 15 October 2021 initially for those with co-morbidities.

As of 07 November 2021, 212 reports were received. Figure 3 shows the distribution of cases by gender and age.




Figure 3. Report distribution by age and gender

Out of the 212 reports, 10 reports were tagged as serious while the rest are non-serious. The most common reported reactions are vaccination/injection site pain, dizziness, headache, blood pressure increased, and tachycardia.

Reports of suspected serious adverse reaction

Adverse reactions experienced after vaccination are considered serious when it resulted to any of the following criteria:

- In-patient hospitalization/prolongation of existing hospitalization
- Significant disability/incapacity
- Life-threatening (e.g. anaphylaxis) and death
- Birth defect or congenital malformations
- Considered to be medically important event

Hypersensitivity including severe allergic reactions

Severe allergic reactions have been reported on the use of COVID-19 vaccines including CoronaVac, COVID-19 Vaccine AstraZeneca, Sputnik V, Comirnaty, Janssen COVID-19 Vaccine, and COVID-19 Vaccine Sinopharm. It only occurs in a few vaccinated individuals. It usually happens in people with a history of severe vaccine reactions. Severe allergic reactions (anaphylaxis) generally occur soon after vaccination and are usually managed with Epinephrine in combination with other medicines. Thus, vaccinees are observed for at least 15 minutes after receiving their vaccine. Epinephrine is readily available in all vaccination sites in case of anaphylaxis.

The proportion of reported side effects of severe allergic reactions to COVID-19 vaccines proved to be statistically rare as the number of vaccinated populations increases. The current reporting rate for anaphylaxis is 4.30 per million doses administered.

Increased blood pressure

Blood pressure (BP) increased has been continuously reported as one of the top adverse reactions to all vaccine platforms. Monitoring BP has been part of the screening process for COVID-19 vaccination program in the country. The program recommends monitoring BP only in vaccine recipients with a history of hypertension, symptomatic hypertension, and based on the clinical judgement of the physician on the vaccination site. This is in relation to the recommendations of [the Philippine Heart Association and Philippine Society of Hypertension on elevated blood pressure readings during COVID-19 vaccination](#).

According to PRESYON 4 (Philippine Heart Association Report on the Study of Hypertension), a nationwide hypertension survey conducted in January to April 2021, the prevalence of hypertension in the Philippines alarmingly increased to 37% in 2021 among adults 18 years old and above from 28% (2013). Out of this 37%, 19% are aware of having hypertension while 18% are unaware. The BP control rate, with or without medications, is 36%. Only about 25% of hypertensive individuals monitor blood pressure at home.³ This study explains the increase in blood pressure observed in most vaccinated individuals.

³ Sison, J.A. (2021, May). Press Conference on PRESYON 4 – Nationwide 2021 Hypertension Survey Results [Video file]. Retrieved from <https://www.facebook.com/philheart.org/videos/159433679504182/>

Thrombosis-thrombocytopenia syndrome

Thrombosis-thrombocytopenia syndrome (TTS) are cases of unusual blood clots with low blood platelets. Following cases of TTS from other countries, COVID-19 Vaccine AstraZeneca and Janssen COVID-19 Vaccine revised their label to include warnings related to thrombosis with thrombocytopenia, a very rare side effect following vaccination.

Vaccinated individuals should watch out for the said adverse event and seek immediate medical assistance if they experience any sign of blood clots and low blood platelet such as: ⁴

- shortness of breath
- chest pain
- leg swelling
- persistent abdominal (belly) pain
- neurological symptoms, such as severe and persistent headaches or blurred vision
- tiny blood spots under the skin beyond the site of the injection

Seven (7) cases of thrombosis have been reported. Causal link of such cases to the vaccination are being reviewed. The FDA together with the Epidemiology Bureau of the DOH shall continue to monitor vaccine safety ensuring that the benefits always outweigh the risks.

⁴ AstraZeneca's COVID-19 vaccine: EMA finds possible link to very rare cases of unusual blood clots with low blood platelets
<https://www.ema.europa.eu/en/news/astrazenecas-covid-19-vaccine-ema-finds-possible-link-very-rare-cases-unusual-blood-clots-low-blood>

Confirmed COVID-19 infections

There were 2,350 confirmed reports of COVID-19 infections. Most of the reported infections were asymptomatic cases. There were 171 severe cases which resulted to a fatal outcome. Most of the fatal reports have not yet completed their vaccination course. Upon assessment, these cases were not related to the use of the vaccine, but these were actual COVID-19 natural infections.

The vaccines currently being used in the COVID-19 vaccination program are non-replicating viral vector, inactivated, and mRNA vaccines. It does not contain any live virus and does not cause COVID-19 infection in vaccine recipients.

Inflammation of the heart

Myocarditis is an inflammation of the heart muscle that may present as chest pain, palpitations, arrhythmias, and/or symptoms of heart failure while pericarditis is an inflammation of the pericardial sac that surrounds the heart and fixes it to the mediastinum. Cases of myocarditis and pericarditis on the use of mRNA vaccine, such as Comirnaty and COVID-19 Vaccine Moderna, have been reported in many countries including the US, UK, Germany, and Israel. Most of the cases are young male. The US FDA announced the revision of fact sheets for Comirnaty and COVID-19 Vaccine Moderna suggesting increased risk of myocarditis and pericarditis following vaccination. EMA's safety committee has also concluded that myocarditis and pericarditis can occur in very rare cases following Comirnaty and COVID-19 Vaccine Moderna.

Vaccinated individuals should watch out for the said adverse event and seek immediate medical assistance if they experience the following symptoms after vaccination:⁵

- breathlessness
- a forceful heartbeat that may be irregular
- chest pain

Six (6) cases of myocarditis and one (1) case of pericarditis have been reported. Causal link of such cases to the vaccination are being reviewed. The FDA together with the Epidemiology Bureau of the DOH shall continue to monitor vaccine safety ensuring that the benefits always outweigh the risks.

⁵ *Comirnaty and Spikevax: possible link to very rare cases of myocarditis and pericarditis*
<https://www.ema.europa.eu/en/news/comirnaty-spikevax-possible-link-very-rare-cases-myocarditis-pericarditis>

Capillary Leak Syndrome

Capillary leak syndrome is a very rare, serious condition that causes fluid leakage from small blood vessels (capillary), resulting in swelling mainly in the arms and legs, low blood pressure, thickening of the blood, and low blood levels of albumin. Several cases were reported on the use of COVID-19 Vaccine AstraZeneca and Janssen COVID-19 Vaccine. The EMA's safety committee recommended contraindication in individuals with previous capillary leak syndrome and inclusion of capillary leak syndrome as a new side effect in the product information for both products.

Vaccinated individuals should watch out for the said adverse event and seek immediate medical assistance if they experience the following symptoms days after vaccination, which may occur together with feeling faint (due to low blood pressure):⁶

- rapid swelling of the arms and legs
- sudden weight gain

No case of capillary leak syndrome has been reported on the use of COVID-19 vaccines in the Philippines as of this time. The FDA together with the Epidemiology Bureau of the DOH shall continue to monitor vaccine safety ensuring that the benefits always outweigh the risks.

⁶ *COVID-19 Vaccine Janssen: Contraindication in individuals with previous capillary leak syndrome and update on thrombosis with thrombocytopenia syndrome*
<https://www.ema.europa.eu/en/medicines/dhpc/covid-19-vaccine-janssen-contraindication-individuals-previous-capillary-leak-syndrome-update>

Guillain-Barré syndrome

Guillain-Barré syndrome (GBS) is a rare, autoimmune disorder in which a person's own immune system damages the nerves, causing muscle weakness and sometimes paralysis. An increased risk for GBS has been observed following vaccination with Janssen COVID-19 Vaccine in the US. The US FDA has announced the revision of fact sheets for Janssen COVID-19 Vaccine to include the observed risk for GBS. EMA's safety committee considered that a causal relationship between Janssen COVID-19 Vaccine and GBS is possible. COVID-19 Vaccine AstraZeneca already updated their product information.

Vaccinated individuals should watch out for the said adverse event and seek immediate medical assistance if they experience signs and symptoms suggestive of GBS such as:⁷

- double vision or difficulty moving eyes
- difficulty swallowing, speaking, or chewing
- coordination problems and unsteadiness
- difficulty walking
- tingling sensations in the hands and feet
- weakness in the limbs, chest, or face
- problems with bladder control and bowel function

Fourteen (14) cases of GBS have been reported. Three (3) cases have been assessed as product related reactions (as per published literature), three (3) cases are indeterminate meaning there is insufficient evidence that the vaccine caused the reaction, and eight (8) are currently being reviewed. The FDA together with the Epidemiology Bureau of the DOH shall continue to monitor vaccine safety ensuring that the benefits always outweigh the risks.

⁷ COVID-19 Vaccine Janssen: Guillain-Barré syndrome listed as a very rare side effect

<https://www.ema.europa.eu/en/news/covid-19-vaccine-janssen-guillain-barre-syndrome-listed-very-rare-side-effect>

Bell's palsy

Bell's palsy is a form of temporary facial paralysis or weakness on one side of the face. It results from dysfunction of facial nerve which directs the muscles on one side of the face. Cases have been reported in a number of people in Hong Kong, Canada, and UK on the use of CoronaVac, Comirnaty, and COVID-19 Vaccine Moderna. The overall number of these reports is relatively small. In relation to this, [CoronaVac vaccination fact sheet was revised to include bell's palsy as a very rare adverse reaction](#) in Hong Kong while Comirnaty product information was revised in Canada. COVID-19 Vaccine Moderna already contains this safety information.

Vaccinated individuals should watch out for the said adverse event and seek immediate medical assistance if they experience any combination of the following symptoms:⁸

- uncoordinated movement of the muscles that control facial expressions, such as smiling, squinting, blinking, or closing the eyelid
- loss of feeling in the face
- headache
- tearing from the eye
- drooling
- lost sense of taste on the front two-thirds of the tongue
- hypersensitivity to sound in the one ear
- inability to close an eye on one side of the face

Twenty-two (22) cases of bell's palsy have been reported. One (1) case have been assessed as product related reactions (as per published literature), four (4) cases are indeterminate meaning there is insufficient evidence that the vaccine caused the reaction, one (1) as coincidental or not related to the vaccine, and (16) are currently being reviewed. The FDA

together with the Epidemiology Bureau of the DOH shall continue to monitor vaccine safety ensuring that the benefits always outweigh the risks.

⁸ *Health Canada updates Pfizer-BioNTech COVID-19 vaccine label to reflect very rare reports of Bell's Palsy*
<https://healthycanadians.gc.ca/recall-alert-rappel-avis/hc-sc/2021/76203a-eng.php>

Cases of hospitalization

One of the criteria for serious adverse reaction is hospitalization or extended hospital stay. Reports of adverse reaction that results in hospitalization does not necessarily mean that vaccine caused the reaction. An Expert Committee reviews and assesses whether the vaccine caused the reaction. Based on the reports received, the hospitalization reporting rate is 3.30 per 100,000 doses administered. Commonly reported causes of hospitalization include pyrexia, cough, dyspnea, and headache.

Reports involving death

As of 07 November 2021, 1,409 fatal events were received. Reports of fatal events does not necessarily mean that the vaccine caused the events. Underlying conditions or pre-existing medical conditions causing fatal events are usually coincidental on the use of the vaccine. It is expected that reports of fatal events will rise as the vaccination program covers more people including those with undiagnosed illness, underlying comorbidities, and pre-existing medical conditions.

The vaccinees reported to have fatal events were aged 18 years and above. The mean age of the fatal cases was 62.5 years. 67.49% (951) of the fatal cases were from age group 60 years and above, 22.43% (316) from age group of 40-59 years, 8.73% (123) from age group 18-39 years, and 1.35 (19) were not identified to what age group they are classified.

Most of these events occurred in persons with multiple existing comorbidities. These include cardiovascular diseases, ischemic heart diseases, cerebrovascular diseases, cancer, diabetes, and infections including pneumonia. There were cases of confirmed COVID-19 infections leading to severe cases with fatal outcomes.

Number of suspected adverse reactions per category

A total of 73,281 case reports consisting of 160,858 suspected adverse reactions were received from the start of the vaccination program. More than one suspected adverse reaction may be reported in a single case. Suspected adverse reactions were coded using the Medical Dictionary for Regulatory Activities (MedDRA) terminology to allow international comparison of reports.

The data presented below are categorized by System Organ Class (SOC), the highest in the hierarchy of MedDRA. They are grouped by manifestation site (e.g. gastrointestinal, cardiac) and etiology (e.g. infections, examinations).

Reactions to inactivated vaccines

- CoronaVac
- COVID-19 Vaccine Sinopharm

Classification	Number of suspected reactions
General symptoms & reactions in the administration site <i>E.g. Pain and reaction in the injection site, chills, discomfort, fever, fatigue</i>	11,988
Cardiac symptoms <i>E.g. Palpitations, bradycardia, tachycardia</i>	860
Ear symptoms <i>E.g. Ear swelling, vertigo, tinnitus, ear discomfort</i>	50
Endocrine symptoms <i>E.g. Adrenal insufficiency, goiter, thyroid symptoms</i>	3
Examinations <i>E.g. Increased blood pressure, increased heart rate, blood glucose increased, SARS-CoV-2 test</i>	11,297
Eye symptoms <i>E.g. Eye itchiness, blurred vision, eye pain, eye swelling</i>	244
Gastrointestinal symptoms <i>E.g. Abdominal pain, diarrhea, nausea, vomiting, dry mouth, lip swelling</i>	3,090
Hepatobiliary symptoms <i>E.g. Jaundice</i>	3
Immune system symptoms <i>E.g. Allergic reactions, hypersensitivity</i>	192
Infections <i>E.g. Cold symptoms, rhinitis</i>	3,038
Metabolism and nutrition-related symptoms <i>E.g. Decreased appetite, increased appetite, starvation, dehydration</i>	402
Musculoskeletal symptoms <i>E.g. Back pain, joint pain, pain in extremities, muscle pain, muscle spasms</i>	1,949
Neoplasm <i>E.g. Liver cancer, endometrial cancer, uterine myoma</i>	3
Neurological symptoms <i>E.g. Dizziness, headache, syncope</i>	7,955
Pregnancy, puerperium, and perinatal conditions <i>E.g. Abortion, hemorrhage</i>	3
Procedural symptoms <i>E.g. Procedural hypertension, vaccination adverse reaction</i>	79
Psychiatric symptoms <i>E.g. Feeling anxious, insomnia, nervousness, disorientation</i>	138
Renal and urinary symptoms <i>E.g. Urine coloring yellow, urine frequency</i>	42
Reproductive symptoms <i>E.g. Vaginal bleeding, vaginal spotting</i>	44
Respiratory symptoms <i>E.g. Cough, nasal congestion, throat irritation</i>	5,575
Skin symptoms <i>E.g. Cold sweat, rash, redness</i>	4,011
Social circumstances <i>E.g. Hearing disability, walking disability</i>	3
Surgical and medical procedure <i>E.g. Tumor debulking, nasolabial flap</i>	1

Symptoms in blood and lymphatic system <i>E.g. Pain in the lymph nodes</i>	35
Vascular symptoms <i>E.g. Flushes, low blood pressure</i>	382




Figure 4. Suspected adverse reaction distribution by SOC for inactivated vaccine

As shown in Figure 4, the SOC which consists of the greatest number of reports were examinations (11,176), followed by general symptoms and reactions in the administration site (8,989), neurological symptoms (6,702), respiratory symptoms (4,223), skin symptoms (3,025), infections (2,659), gastrointestinal symptoms (2,327), musculoskeletal symptoms (1,708), cardiac symptoms (833), and metabolism and nutrition-related symptoms (400).

The top reported events for CoronaVac are:

- blood pressure increased (37.99%)
- headache (13.08%)
- pyrexia (13.05%)
- vaccination/injection site pain (10.25%)
- cough (9.36%)
- rash (7.00%)
- dizziness (6.63%)
- nasopharyngitis (5.57%)
- malaise (5.03%)
- dyspnea (4.41%)

The top reported events for COVID-19 Vaccine Sinopharm are:

- pyrexia (28.44%)
- cough (16.51%)
- dizziness (14.68%)
- headache (13.76%)
- rash (11.01%)
- nasopharyngitis (9.17%),
- dyspnea (8.26%), blood pressure increased (8.26%)
- fatigue (7.34%), pruritus (7.34%)

Reactions to non-replicating viral vector vaccines

- COVID-19 vaccine AstraZeneca
- Sputnik V
- Janssen COVID-19 Vaccine

Classification	Number of suspected reactions
General symptoms & reactions in the administration site <i>E.g. Pain and reaction in the injection site, chills, discomfort, fever, fatigue</i>	44,024
Cardiac symptoms <i>E.g. Palpitations, bradycardia, tachycardia</i>	741
Congenital, familial, and genetic disorder <i>E.g. Polycystic kidney</i>	1
Ear symptoms <i>E.g. Ear swelling, vertigo, tinnitus, ear discomfort</i>	62
Endocrine symptoms <i>E.g. Adrenal insufficiency, goiter, thyroid symptoms</i>	3
Examinations <i>E.g. Increased blood pressure, increased heart rate, blood glucose increased, SARS-CoV-2 test</i>	6,656
Eye symptoms <i>E.g. Eye itchiness, blurred vision, eye pain, eye swelling</i>	482
Gastrointestinal symptoms <i>E.g. Abdominal pain, diarrhea, nausea, vomiting, lip swelling</i>	4,709
Hepatobiliary symptoms <i>E.g. Jaundice</i>	5
Immune system symptoms <i>E.g. Allergic reactions, hypersensitivity</i>	271
Infections <i>E.g. Cold symptoms, rhinitis</i>	2,663
Metabolism and nutrition-related symptoms <i>E.g. Decreased appetite, increased appetite, starvation, dehydration</i>	728
Musculoskeletal symptoms <i>E.g. Back pain, joint pain, pain in extremities, muscle pain, muscle spasms</i>	8,990
Neoplasm <i>E.g. Liver cancer, endometrial cancer, uterine myoma</i>	2
Neurological symptoms <i>E.g. Dizziness, headache, syncope</i>	16,340
Pregnancy, puerperium, and perinatal conditions <i>E.g. Abortion, hemorrhage</i>	9

Procedural symptoms <i>E.g. Procedural hypertension, vaccination adverse reaction</i>	156
Psychiatric symptoms <i>E.g. Feeling anxious, insomnia, nervousness, disorientation</i>	105
Renal and urinary symptoms <i>E.g. Urine coloring yellow, urine frequency</i>	40
Reproductive symptoms <i>E.g. Vaginal bleeding, vaginal spotting</i>	55
Respiratory symptoms <i>E.g. Cough, nasal congestion, throat irritation</i>	4,169
Skin symptoms <i>E.g. Cold sweat, rash, redness</i>	3,441
Social circumstances <i>E.g. Hearing disability, walking disability</i>	2
Surgical and medical procedure <i>E.g. Tumor debulking, nasolabial flap</i>	2
Symptoms in blood and lymphatic system <i>E.g. Pain in the lymph nodes</i>	68
Vascular symptoms <i>E.g. Flushes, low blood pressure</i>	372




Figure 5. Suspected adverse reaction distribution by SOC for viral vector vaccines

As shown in Figure 5, the SOC which consists of the greatest number of reports were general symptoms and reactions in the administration site (23,098), followed by neurological

symptoms (14,236), musculoskeletal symptoms (7,282), examinations (6,614), gastrointestinal symptoms (3,608), respiratory symptoms (3,328), skin symptoms (2,659), infections (2,351), metabolism and nutrition-related symptom (721), and cardiac symptoms (709).

The top reported events for COVID-19 Vaccine AstraZeneca are:

- pyrexia (40.46%)
- headache (34.09%)
- vaccination/injection site pain (23.44%)
- malaise (21.72%)
- chills (16.46%)
- myalgia (15.81%)
- blood pressure increased (15.52%)
- fatigue (11.90%)
- arthralgia (7.81%)
- dizziness (6.38%)

The top reported events for Sputnik V are:

- blood pressure increased (57.20%)
- pyrexia (11.44%)
- headache (8.48%)
- cough (6.33%)
- rash (5.25%)
- nasopharyngitis (4.85%)
- heart rate increased (4.71%)
- vaccination/injection site pain (4.44%)
- dizziness (4.17%)
- COVID-19 (3.50%)

The top reported events for Janssen COVID-19 Vaccine are:

- blood pressure increased (27.26%)
- pyrexia (20.36%)
- vaccination/injection site pain (17.63%)
- headache (11.50%)
- cough (10.93%)
- dyspnea (6.04%)
- hypoesthesia (5.81%)
- malaise (5.58%)
- dizziness (4.86%)
- rash (4.80%)

Reactions to mRNA vaccines

- Comirnaty
- COVID-19 Vaccine Moderna

Classification	Number of suspected reactions
General symptoms & reactions in the administration site <i>E.g. Pain and reaction in the injection site, chills, discomfort, fever, fatigue</i>	4,924
Cardiac symptoms <i>E.g. Palpitations, bradycardia, tachycardia</i>	385
Ear symptoms <i>E.g. Ear swelling, vertigo, tinnitus, ear discomfort</i>	19
Examinations <i>E.g. Increased blood pressure, increased heart rate, blood glucose increased, SARS-CoV-2 test</i>	2,469
Eye symptoms <i>E.g. Eye itchiness, blurred vision, eye pain, eye swelling</i>	94
Gastrointestinal symptoms <i>E.g. Abdominal pain, diarrhea, nausea, vomiting, lip swelling</i>	807
Immune system symptoms <i>E.g. Allergic reactions, hypersensitivity</i>	64
Infections <i>E.g. Cold symptoms, rhinitis</i>	624
Metabolism and nutrition-related symptoms <i>E.g. Decreased appetite, increased appetite, starvation, dehydration</i>	108
Musculoskeletal symptoms <i>E.g. Back pain, joint pain, pain in extremities, muscle pain, muscle spasms</i>	837
Neoplasm <i>E.g. Breast cancer</i>	1
Neurological symptoms <i>E.g. Dizziness, headache, syncope</i>	2,296
Pregnancy, puerperium, and perinatal conditions <i>E.g. Abortion, hemorrhage</i>	4
Procedural symptoms <i>E.g. Procedural hypertension, vaccination adverse reaction</i>	30
Psychiatric symptoms <i>E.g. Feeling anxious, insomnia, nervousness, disorientation</i>	55
Renal and urinary symptoms <i>E.g. Urine coloring yellow, urine frequency</i>	18
Reproductive symptoms <i>E.g. Vaginal bleeding, vaginal spotting</i>	30
Respiratory symptoms <i>E.g. Cough, nasal congestion, throat irritation</i>	1,344
Skin symptoms <i>E.g. Cold sweat, rash, redness</i>	1,078
Social circumstances <i>E.g. Hearing disability, walking disability</i>	2
Symptoms in blood and lymphatic system <i>E.g. Pain in the lymph nodes</i>	48
Surgical and medical procedure <i>E.g. Chemotherapy</i>	1
Vascular symptoms <i>E.g. Flashes, low blood pressure</i>	105




Figure 6. Suspected adverse reaction distribution by SOC for mRNA vaccine

As shown in Figure 6, the SOC which consists of the greatest number of reports were general symptoms and reactions in the administration site (3,059), followed by examinations (2,421), neurological symptoms (1,891), respiratory symptoms (1,037), skin symptoms (822), musculoskeletal symptoms (683), gastrointestinal symptoms (588), infections (562), cardiac symptoms (380), and metabolism and nutrition-related symptoms (108).

The top reported events for Comirnaty are:

- blood pressure increased (33.56%)
- pyrexia (17.66%)
- headache (12.37%)
- vaccination/injection site pain (9.19%)
- cough (8.39)
- dizziness (7.88%)
- rash (6.89%)
- nasopharyngitis (4.87%)
- dyspnea (4.72%)
- malaise (4.57%)

The top reported events for COVID-19 vaccine Moderna are:

- pyrexia (31.80%)
- vaccination/injection site pain (24.37%)
- blood pressure increased (21.08%)
- headache (19.93%)

- chills (9.60%)
- cough (8.62%)
- malaise (7.38%)
- dizziness (6.83%)
- rash (6.74%)
- pain (6.53%)

Outcome of suspected adverse reactions

The outcome of cases of suspected adverse reactions to COVID-19 vaccines is shown in Figure 7. Overall, most of the reported cases have *recovered/resolved* (77.36%), although there were few cases who have *recovered but with sequelae* (0.03%). Over 12% of the cases are *recovering/resolving* while less than 1% have *not recovered/not resolved* at the time of reporting. A proportion of 1.92% were reported with fatal outcomes as discussed in the section reports involving death.




Figure 7. Case outcome

Reporting of suspected adverse reactions following vaccination

Individuals who have received their COVID-19 vaccination shots can report any suspected adverse reaction to any of the following:

- Immunization site where you were vaccinated
- Directly to the vaccine manufacturer or emergency use authorization holder
 - [Sinovac – CoronaVac](#)
 - [AstraZeneca – COVID-19 Vaccine AstraZeneca](#)
 - [Gamaleya – Sputnik V](#)
 - [Pfizer – Comirnaty](#)
 - [Zuellig – COVID-19 Vaccine Moderna](#)
 - Johnson & Johnson – Janssen COVID-19 Vaccine
- [FDA online reporting system](#)

Kindly **report only to one** of the above to avoid duplication of reports.