



Reports of Suspected Adverse Reaction to COVID-19 Vaccines (01 March 2021 to 27 February 2022)

Contents

About the report	2
Summary	3
COVID-19 vaccines with Emergency Use Authorization in the Philippines	4
Statistics regarding reports of suspected adverse reactions	4
Distribution of reports of adverse reaction for each vaccine.....	5
Demographics	5
Pregnant women and lactating mothers.....	6
Vaccination in adolescent population	7
Booster shots	7
National Vaccination Days	8
Roll-out in pharmacies and clinics	8
Resbakuna kids	8
Reports of suspected serious adverse reaction	9
Number of suspected adverse reactions per category	14
Reactions to inactivated vaccines	14
Reactions to non-replicating viral vector vaccines	17
Reactions to mRNA vaccines	20
Outcome of suspected adverse reaction	22
Reporting of suspected adverse reactions following vaccination	23



About the report

- A summary is presented below of all received suspected adverse reaction reports following COVID-19 vaccination from 01 March 2021, the date when the first vaccine became available, up to 27 February 2022.
- Eight (8) vaccines under Emergency Use Authorization (EUA) are currently being used in the vaccination program: CoronaVac, COVID-19 Vaccine AstraZeneca, Sputnik V, Comirnaty, COVID-19 Vaccine Moderna, Janssen COVID-19 Vaccine, COVID-19 Vaccine Sinopharm, and Sputnik Light.
- Data are based on the VigiFlow, the national database of adverse reactions in the Philippines. It includes reports from various epidemiology surveillance units (ESUs) of the Department of Health (DOH), vaccination sites, hospitals, patients/consumers, and EUA holders.
- Symptoms or diseases that occur after vaccination are reported if there is a *suspicion* of a possible link. However, it cannot be assumed that there is a causal relationship between the suspected adverse reaction and the vaccine.
- This report contains all suspected adverse reactions regardless of any possible causal relationship.
- Additional information may become available in individual case reports at any time, which may change the assessment and figures presented.
- Adverse reaction reports are necessary for the safety assessment of the vaccines, making sure that the benefits always outweigh the risks.
- Reports are constantly reviewed and monitored for the possible emergence/identification of unknown adverse reactions also known as signal. If a signal is identified, the FDA and DOH perform investigations, regulatory actions, and timely communication.
- A weekly report is published to summarize reported adverse reactions to the COVID-19 vaccines.

Summary

This report is based on an assessment of adverse reaction reports received by 27 February 2022. As per benefit-risk assessment, these reports do not provide a basis for revising the current recommendations regarding the use of COVID-19 vaccines.

A report of adverse reaction does not necessarily mean that the vaccine caused the reactions. A mere suspicion may also be reported. Undiagnosed illness, underlying comorbidities, and pre-existing medical conditions unrelated to vaccination can be factors in reporting adverse reactions. The relative numbers should not be used to compare the safety of different vaccines.

Like any other vaccines, COVID-19 vaccines may cause adverse reactions in some people while others may not experience any adverse reaction. It is possible for several persons to experience the same adverse event but for the report to be serious for one person and non-serious for another person. Most of the reported reactions are generally in line with what is described in the product information and labels. Such reports are minor adverse reactions, which include body pain, chills, fatigue, fever, headache, nausea, and pain in the injection site. These usually appear on the first or second day of vaccination and may last for 2-3 days. Most people tolerate these adverse reactions while others experience greater discomfort.

Serious adverse reactions have also been reported. The FDA together with other public health partners are continuously monitoring the adverse experience as more people are being vaccinated with COVID-19 vaccines. Such monitoring will provide reassurance that the vaccines are safe and effective for use.

Considering the post-authorization experience on the use of COVID-19 Vaccine AstraZeneca and Janssen COVID-19 Vaccine of other countries, information on the very rare and serious adverse events of thrombosis and thrombocytopenia and in some cases accompanied by bleeding have been revised under the special warning and precautions for use in their product labels. In addition, capillary leak syndrome and Guillain-Barré syndrome were added as a rare new side effect.

The label of mRNA vaccines Moderna and Comirnaty (Pfizer-BioNTech) have been revised to include imposition of the European Medicines Agency and the USFDA to include safety information on myocarditis and pericarditis.

The immunization program expanded its coverage to include adolescent individuals (12 to 17 years old). Pfizer and Moderna are the only vaccines with EUA for the said population. The rollout started on the second week of October. To further expand vaccine coverage, vaccination to children ages 5-11 years started on February 2022. Likewise, monitoring of suspected adverse reactions was extended to include such population.

Eight (8) vaccines are currently used in the immunization program. Supplies of vaccines are either procured by the government and/or private sector or supplied under the COVAX facility.

COVID-19 vaccines with Emergency Use Authorization in the Philippines

At present, the FDA granted [nine \(9\) COVID-19 vaccines with emergency use authorization](#):

- Pfizer-BioNTech COVID-19 Vaccine (Comirnaty)
- ChAdOx1-S [recombinant] (COVID-19 Vaccine AstraZeneca)
- SARS-CoV-2 Vaccine (Vero Cell), Inactivated (CoronaVac)
- Gam-COVID-Vac (Sputnik V & Sputnik Light)
- Ad26.COV2-S [recombinant] (Janssen COVID-19 Vaccine)
- Whole Virion Inactivated Corona Virus (Covaxin)
- COVID-19 mRNA Vaccine [nucleoside modified] (COVID-19 Vaccine Moderna)
- Inactivated COVID-19 Vaccine (Vero Cell) (COVID-19 Vaccine Sinopharm BIBP/Wuhan)
- SARS-CoV-2 rS Protein Nanoparticle Vaccine [Covovax]

Various vaccine platforms have been approved for use in the Philippines. Comirnaty and COVID-19 Vaccine Moderna are mRNA vaccines; COVID-19 Vaccine AstraZeneca and Janssen COVID-19 Vaccine are non-replicating viral vector vaccines while Sputnik V uses the same technology having two (2) different (dose) components of viral vectors, and Spuntik Light having a single dose component; CoronaVac, Covaxin, and COVID-19 Vaccine Sinopharm BIBP/Wuhan are inactivated vaccines; and Covovax is a protein subunit vaccine. All are administered in two doses within an interval of a few weeks except for Janssen COVID-19 Vaccine and Sputnik Light, which are administered as a single-dose.

Statistics regarding reports of suspected adverse reactions

As of 27 February 2022, more than 135.4 million doses of COVID-19 vaccines (either CoronaVac, COVID-19 Vaccine AstraZeneca, Sputnik V, Comirnaty, COVID-19 Vaccine Moderna, Janssen COVID-19 Vaccine, COVID-19 Vaccine Sinopharm, or Sputnik Light) were administered. Over 63 million individuals are now fully vaccinated (either given a single-dose or 2-dose vaccine course) while more than 5.6 million are partly vaccinated waiting for their second dose to be administered. 10,137,875 million individuals already received their third or additional doses (either homologous or heterologous booster doses). A total of 92,215 suspected adverse reaction reports were received, evaluated, and analyzed by the FDA. To disaggregate, 33,387 have been reported for CoronaVac, 35,032 for COVID-19 Vaccine AstraZeneca, 814 for Sputnik V, 13,612 for Comirnaty, 4,954 for COVID-19 Vaccine Moderna, 4,205 for Janssen COVID-19 Vaccine, and 211 for COVID-19 Vaccine Sinopharm.

Table 1. Data on vaccination and suspected adverse reaction reports.

Indicators	Value
Total number of doses administered	135,479,183
No. of fully vaccinated individuals	63,092,684
No. of individuals partly vaccinated	5,637,883
No. of individuals with booster shots	10,137,875

No. of suspected adverse reaction reports	92,215 (0.07% of doses administered)
No. of suspected serious adverse reaction reports	5,932 (0.004% of doses administered)

Distribution of reports of adverse reactions for each vaccine

Data shown below are cumulative reports from the start of the vaccination program on 01 March 2021 up until 20 February 2022.

Table 2. Distribution of reports of adverse reactions for each vaccine

Vaccine	Date started	Total vaccine doses administered ^b	Number of fully vaccinated individuals ^b	Number of individuals partly Vaccinated	Number of individuals with booster shots	Total number of reports ^a	Reports of non-serious events	Reports of serious events
CoronaVac	01 Mar 2021	44,530,727	21,316,147	1,295,558	602,875	33,387	30,902	2,485
AstraZeneca	07 Mar 2021	20,362,040	8,415,903	1,232,993	2,297,241	35,032	33,694	1,338
Sputnik V	04 May 2021	1,080,022	512,176	55,441	229	814	782	32
Comirnaty	13 May 2021	45,174,730	19,091,517	2,254,600	4,737,096	13,612	12,598	1,014
Moderna	30 June 2021	16,817,400	6,787,506	743,867	2,498,521	4,954	4,568	386
Janssen	20 July 2021	6,482,327	6,481,106	-	1,221	4,205	3,563	642
Sinopharm	25 Aug 2021	1,030,846	487,492	55,424	438	211	176	35
Sputnik Light	2022	1,091	837	-	254	0	0	0
TOTAL		135,479,183	63,092,684	5,637,883	10,137,875	92,215	86,283	5,932

Data source: ^aVigiFlow, ^bNVOC daily report as 6PM, 27 February 2022

Notes: Additional information may become available in individual cases, which may change the figures presented

^aAn individual is considered partly vaccinated if they have received only one dose of a two-dose vaccine course. An individual is considered fully vaccinated if they have received a single-dose vaccine or both doses of a two-dose vaccine

^cData concerning various vaccines are not directly comparable. COVID-19 vaccines profile varies, they have not been used for equal periods of time and they have been administered to number of people with different profiles including various age and sex.

Demographics

The figures below provide a descriptive overview of the population who have experienced adverse reactions to COVID-19 vaccines. Figure 1 and Figure 2 shows the distribution of reports by gender and age. Click [here](#) to show disaggregated data.




Figure 1. Report distribution by gender




Figure 2. Report distribution by age

The early phase of the vaccination program is intended for the frontline health workers, thus, the high number in the female and younger population might be associated with the data that our health system is dominated by female (75%) and young adults under the age of 35 (65%).¹ An increasing number of reports from the age group 40 years and above have been observed in the past few weeks of the vaccination program. This may be attributed to the coverage of priority groups of senior citizens and individuals with comorbidities.

Relative to the inclusion of the frontline personnel in the priority groups, the observed increasing number of reports in the male population may be attributed to the vaccine coverage and statistics that more males are employed than females (6 in every 10).²

¹ Human Resource for Health in the Time of the COVID-19 Pandemic: Does the Philippines Have Enough? <https://www.drdf.org.ph/sites/default/files/pdf/COVID-19-Research-Brief-08.pdf>

² Employment situation in July 2018, Philippine Statistics Authority <https://psa.gov.ph/statistics/survey/labor-and-employment/labor-force-survey/title/Employment%20Situation%20in%20July%202018>

Pregnant women and lactating mothers

Pregnant women are included in the A3 priority group for COVID-19 vaccination. **Pregnancy is not a contraindication to getting the COVID-19 vaccine (except for the Gamaleya vaccine, Gamaleya vaccine shall not be administered to the pregnant and breastfeeding populations).** They are considered to have a low risk of contracting COVID-19; however, studies have shown that pregnant women have a higher risk of having severe COVID-19 infection compared to non-pregnant women thus the benefits of getting vaccinated outweigh the risks.

As of 27 February 2022, 215 suspected adverse reaction reports were received from pregnant women. Out of these reports, 105 reports were tagged as serious and 110 reports were non-serious. The most commonly reported adverse reactions from pregnant women include vaccination/injection site pain, headache, pyrexia, COVID-19, cough, dyspnea and nasopharyngitis.

Breastfeeding is vital to the health of infants and their mothers. COVID-19 vaccination is also recommended for breastfeeding mothers. Limited data is available on the effects of the

vaccine on milk production and excretion. As of 27 February 2022, 166 reports were received from the group of lactating mothers. Nine (9) reports were tagged as serious and 157 were non-serious. The most commonly reported adverse reactions from lactating mothers include pyrexia, headache, vaccination/injection site pain, malaise, and myalgia.

Overall, data suggests that the benefits of receiving a COVID-19 vaccine outweigh any known or potential risks of vaccination during pregnancy and lactation.

References:

WHO COVID-19 vaccines and pregnancy, breastfeeding, fertility

CDC COVID-19 Vaccines While Pregnant or Breastfeeding

Vaccination in adolescent population

The roll out for vaccinating adolescent population (12-17 years old) started last 15 October 2021 initially for those with co-morbidities and expanded to include all adolescents (with or without co-morbidities) on 02 November 2021. Pfizer and Moderna are the only vaccines with EUA for the adolescent population.

As of 27 February 2022, 3,138 reports were received. One hundred ninety-one (191) reports were tagged as serious (145 hospitalizations) and 2,898 reports are non-serious. The most common reported reactions are dizziness, pyrexia, vaccination/injection site pain, headache, and blood pressure increased.

Although very rare cases of myocarditis and pericarditis have been observed in this population, the benefits of using such vaccines still outweigh the risks. There are two (2) cases of myocarditis and one (1) case of pericarditis that have been reported in adolescent population. One (1) case of myocarditis has been assessed as product related reactions (as per published literature) while one (1) case of myocarditis including one (1) case of pericarditis are currently being reviewed.

Booster shots

The roll out of vaccination for the booster shots (third or additional doses) started last 17 November 2021 initially for healthcare workers. Senior citizens, immunocompromised, and individuals with comorbidities at high risk of developing severe COVID-19 on 22 November 2021 followed this. The general population ages 18 years and above who completed their vaccine doses may also receive booster shots, which started last 03 December 2021.

Individuals eligible for booster shots have the option of receiving a [homologous or heterologous booster](#) dose given that they have completed their primary dose series (at least six (6) months after getting the second dose of either CoronaVac, COVID-19 Vaccine AstraZeneca, Sputnik V, Comirnaty, or COVID-19 Vaccine Moderna and at least 3 months for Janssen COVID-19 vaccine). Last 21 December 2021, [interval for the administration of booster doses was shortened](#) to at least three (3) months after the 2nd dose of a two-dose vaccine while at least two (2) months for a single-dose vaccine.

As of 27 February 2022, 2,767 suspected adverse reaction reports were received, 158 reports were tagged as serious while 2,554 are tagged as non-serious, and 55 reports with no tag whether serious or non-serious. The most common reported adverse reaction includes vaccination/injection site pain, pyrexia, headache, malaise, and blood pressure increased.

National Vaccination Days

The Bayanihan, Bakunahan National Vaccination Days were held last 29 November to 02 December 2021. A total of 1,768 reports were received out of more than 8 million doses administered. Twenty-six (26) serious adverse reaction were reported while the rest are non-serious. The top reported reactions were blood pressure increased, dizziness, headache, pyrexia, and vaccination/injection site pain.

With the aim of increasing COVID-19 vaccination coverage nationwide and supporting the national government's goal of giving Filipino a safe and healthy Christmas, another round of the National Vaccination Days was organized from 15 December to 17 December 2021 and extended until 22 December due to typhoon "Odette". Unaffected areas in certain parts of Luzon initiated the implementation of the National Vaccination Days 2, while areas in the Visayas and Mindanao will follow. As of 19 December 2021, a total of 177 reports were received out of 2.7 million doses administered. Most of the reported cases involves minor adverse reactions. The top reported reactions were blood pressure increased, vaccination/injection site pain, dizziness, headache, and vomiting.

The third wave of Bayanihan, Bakunahan National Vaccination Days started last 10 February 2022 and will run until 18 February 2022. As of 18 February 2022, 168 reports were received. Five (5) reports were tagged as serious, 161 reports as non-serious, and two (2) reports with no tag whether serious or non-serious. The top reported reactions were dizziness, pyrexia, blood pressure increased, rash and vaccination/injection site pain.

Roll out in pharmacies and clinics

The COVID-19 vaccination program aims to reach more Filipinos through its vaccination roll out in pharmacies and clinics which started last 20 January 2022 initially for booster shots. High-risk groups and children below 18 years old are not qualified to be vaccinated in drugstores while clinics may give shots to high-risk group such as senior citizens and individuals with co-morbidities. Initially, CoronaVac and COVID-19 Vaccine AstraZeneca are the only vaccines being utilized in these settings.

Resbakuna kids

Resbakuna kids started last 07 February 2022. This is to offer additional protection for children ages 5-11 years old.

As of 27 February 2022, 239 reports were received. Four (4) reports were tagged as serious and 235 reports are non-serious. The most common reported reactions are vaccination/injection site pain, rash pyrexia, vomiting, and pruritus.

Reports of suspected serious adverse reaction

Adverse reactions experienced after vaccination are considered serious when it resulted to any of the following criteria:

- In-patient hospitalization/prolongation of existing hospitalization
- Significant disability/incapacity
- Life-threatening (e.g. anaphylaxis) and death
- Birth defect or congenital malformations
- Considered to be medically important event

Hypersensitivity including severe allergic reactions

Severe allergic reactions have been reported on the use of COVID-19 vaccines including CoronaVac, COVID-19 Vaccine AstraZeneca, Sputnik V, Comirnaty, COVID-19 Vaccine Moderna, Janssen COVID-19 Vaccine, and COVID-19 Vaccine Sinopharm. It only occurs in a few vaccinated individuals. It usually happens in people with a history of severe vaccine reactions. Severe allergic reactions (anaphylaxis) generally occur soon after vaccination and are usually managed with Epinephrine in combination with other medicines. Thus, vaccinees are observed for at least 15 minutes after receiving their vaccine. Epinephrine is readily available in all vaccination sites in case of anaphylaxis.

The proportion of reported side effects of severe allergic reactions to COVID-19 vaccines proved to be statistically rare as the number of vaccinated populations increases. The current reporting rate for anaphylaxis is 2.30 per million doses administered.

Increased blood pressure

Blood pressure (BP) increased has been continuously reported as one of the top adverse reactions to all vaccine platforms. Monitoring BP has been part of the screening process for COVID-19 vaccination program in the country. The program recommends monitoring BP only in vaccine recipients with a history of hypertension, symptomatic hypertension, and based on the clinical judgement of the physician on the vaccination site. This is in relation to the recommendations of [the Philippine Heart Association and Philippine Society of Hypertension on elevated blood pressure readings during COVID-19 vaccination](#).

According to PRESYON 4 (Philippine Heart Association Report on the Study of Hypertension), a nationwide hypertension survey conducted in January to April 2021, the prevalence of hypertension in the Philippines alarmingly increased to 37% in 2021 among adults 18 years old and above from 28% (2013). Out of this 37%, 19% are aware of having hypertension while 18% are unaware. The BP control rate, with or without medications, is 36%. Only about 25% of hypertensive individuals monitor blood pressure at home.³ This study explains the increase in blood pressure observed in most vaccinated individuals.

³ Sison, J.A. (2021, May). Press Conference on PRESYON 4 – Nationwide 2021 Hypertension Survey Results [Video file]. Retrieved from <https://www.facebook.com/philheart.org/videos/159433679504182/>

Thrombosis-thrombocytopenia syndrome

Thrombosis-thrombocytopenia syndrome (TTS) are cases of unusual blood clots with low blood platelets. Following cases of TTS from other countries, COVID-19 Vaccine AstraZeneca and Janssen COVID-19 Vaccine revised their label to include warnings related to thrombosis with thrombocytopenia, a very rare side effect following vaccination.

Vaccinated individuals should watch out for the said adverse event and seek immediate medical assistance if they experience any sign of blood clots and low blood platelet such as: ⁴

- shortness of breath
- chest pain
- leg swelling
- persistent abdominal (belly) pain
- neurological symptoms, such as severe and persistent headaches or blurred vision
- tiny blood spots under the skin beyond the site of the injection

Ten (10) cases of thrombosis have been reported. One (1) case has been assessed as indeterminate meaning there is insufficient evidence that the vaccine caused the reaction and nine (9) cases are currently being reviewed. Causal link of such cases to the vaccination are being reviewed. The FDA together with the Epidemiology Bureau of the DOH shall continue to monitor vaccine safety ensuring that the benefits always outweigh the risks.

⁴ AstraZeneca's COVID-19 vaccine: EMA finds possible link to very rare cases of unusual blood clots with low blood platelets
<https://www.ema.europa.eu/en/news/astrazenecas-covid-19-vaccine-ema-finds-possible-link-very-rare-cases-unusual-blood-clots-low-blood>

Confirmed COVID-19 infections

There were 3,552 confirmed reports of COVID-19 infections. Most of the reported infections were asymptomatic cases. There were 221 severe cases that resulted to a fatal outcome. Most of the fatal reports have not yet completed their vaccination course. Upon assessment, these cases were not related to the use of the vaccine, but these were actual COVID-19 natural infections.

The vaccines currently being used in the COVID-19 vaccination program are non-replicating viral vector, inactivated, and mRNA vaccines. It does not contain any live virus and therefore does not cause COVID-19 infection in vaccine recipients.

Inflammation of the heart

Myocarditis is an inflammation of the heart muscle that may present as chest pain, palpitations, arrhythmias, and/or symptoms of heart failure while pericarditis is an inflammation of the pericardial sac that surrounds the heart and fixes it to the mediastinum. Cases of myocarditis and pericarditis on the use of mRNA vaccine, such as Comirnaty and COVID-19 Vaccine Moderna, have been reported in many countries including the US, UK, Germany, and Israel. Most of the cases are young male. The US FDA announced the revision of fact sheets for Comirnaty and COVID-19 Vaccine Moderna suggesting increased risk of myocarditis and pericarditis following vaccination. EMA's safety committee has also concluded that myocarditis and pericarditis can occur in very rare cases following Comirnaty and COVID-19 Vaccine Moderna.

Vaccinated individuals should watch out for the said adverse event and seek immediate medical assistance if they experience the following symptoms after vaccination:⁵

- breathlessness
- a forceful heartbeat that may be irregular
- chest pain

Nine (9) cases of myocarditis and two (2) cases of pericarditis have been reported. Two (2) cases of myocarditis have been assessed as product related reactions (as per published literature) and seven (7) cases including two (2) cases of pericarditis are currently being reviewed. Causal link of such cases to the vaccination are being reviewed. The FDA together with the Epidemiology Bureau of the DOH shall continue to monitor vaccine safety ensuring that the benefits always outweigh the risks.

⁵ *Comirnaty and Spikevax: possible link to very rare cases of myocarditis and pericarditis*
<https://www.ema.europa.eu/en/news/comirnaty-spikevax-possible-link-very-rare-cases-myocarditis-pericarditis>

Capillary Leak Syndrome

Capillary leak syndrome is a very rare, serious condition that causes fluid leakage from small blood vessels (capillary), resulting in swelling mainly in the arms and legs, low blood pressure, thickening of the blood, and low blood levels of albumin. Several cases were reported on the use of COVID-19 Vaccine AstraZeneca and Janssen COVID-19 Vaccine. The EMA's safety committee recommended contraindication in individuals with previous capillary leak syndrome and inclusion of capillary leak syndrome as a new side effect in the product information for both products.

Vaccinated individuals should watch out for the said adverse event and seek immediate medical assistance if they experience the following symptoms days after vaccination, which may occur together with feeling faint (due to low blood pressure):⁶

- rapid swelling of the arms and legs
- sudden weight gain

No case of capillary leak syndrome has been reported on the use of COVID-19 vaccines in the Philippines as of this time. The FDA together with the Epidemiology Bureau of the DOH shall continue to monitor vaccine safety ensuring that the benefits always outweigh the risks.

⁶ *COVID-19 Vaccine Janssen: Contraindication in individuals with previous capillary leak syndrome and update on thrombosis with thrombocytopenia syndrome*
<https://www.ema.europa.eu/en/medicines/dhpc/covid-19-vaccine-janssen-contraindication-individuals-previous-capillary-leak-syndrome-update>

Guillain-Barré syndrome

Guillain-Barré syndrome (GBS) is a rare, autoimmune disorder in which a person's own immune system damages the nerves, causing muscle weakness and sometimes paralysis. An increased risk for GBS has been observed following vaccination with Janssen COVID-19 Vaccine in the US. The US FDA has announced the revision of fact sheets for Janssen COVID-

19 Vaccine to include the observed risk for GBS. EMA's safety committee considered that a causal relationship between Janssen COVID-19 Vaccine and GBS is possible. COVID-19 Vaccine AstraZeneca already updated their product information.

Vaccinated individuals should watch out for the said adverse event and seek immediate medical assistance if they experience signs and symptoms suggestive of GBS such as:⁷

- double vision or difficulty moving eyes
- difficulty swallowing, speaking, or chewing
- coordination problems and unsteadiness
- difficulty walking
- tingling sensations in the hands and feet
- weakness in the limbs, chest, or face
- problems with bladder control and bowel function

Twenty (20) cases of GBS have been reported. Four (4) cases have been assessed as product related reactions (as per published literature), four (4) cases are indeterminate meaning there is insufficient evidence that the vaccine caused the reaction, and 12 are currently being reviewed. The FDA together with the Epidemiology Bureau of the DOH shall continue to monitor vaccine safety ensuring that the benefits always outweigh the risks.

⁷ COVID-19 Vaccine Janssen: Guillain-Barré syndrome listed as a very rare side effect
<https://www.ema.europa.eu/en/news/covid-19-vaccine-janssen-guillain-barre-syndrome-listed-very-rare-side-effect>

Bell's palsy

Bell's palsy is a form of temporary facial paralysis or weakness on one side of the face. It results from dysfunction of facial nerve, which directs the muscles on one side of the face. Cases have been reported in a number of people in Hong Kong, Canada, and UK on the use of CoronaVac, Comirnaty, and COVID-19 Vaccine Moderna. The overall number of these reports is relatively small. In relation to this, [CoronaVac vaccination fact sheet was revised to include bell's palsy as a very rare adverse reaction](#) in Hong Kong while Comirnaty product information was revised in Canada. COVID-19 Vaccine Moderna already contains this safety information.

Vaccinated individuals should watch out for the said adverse event and seek immediate medical assistance if they experience any combination of the following symptoms:⁸

- uncoordinated movement of the muscles that control facial expressions, such as smiling, squinting, blinking, or closing the eyelid
- loss of feeling in the face
- headache
- tearing from the eye
- drooling
- lost sense of taste on the front two-thirds of the tongue
- hypersensitivity to sound in the one ear
- inability to close an eye on one side of the face

Twenty-five (25) cases of Bell's palsy have been reported. Three (3) cases have been assessed as product related reactions (as per published literature), four (4) cases are indeterminate meaning there is insufficient evidence that the vaccine caused the reaction, two (2) cases as coincidental or not related to the vaccine, and 16 cases are currently being reviewed. The FDA together with the Epidemiology Bureau of the DOH shall continue to monitor vaccine safety ensuring that the benefits always outweigh the risks.

⁸ Health Canada updates Pfizer-BioNTech COVID-19 vaccine label to reflect very rare reports of Bell's Palsy
<https://healthycanadians.gc.ca/recall-alert-rappel-avis/hc-sc/2021/76203a-eng.php>

Immune thrombocytopenia

Immune thrombocytopenia is an autoimmune condition in which the immune system mistakenly targets blood cells called platelets that are needed for normal blood clotting. Very rare cases have been reported internationally after receiving the Janssen COVID-19 Vaccine and the COVID-19 Vaccine AstraZeneca. The product information for both vaccines have been recommended to update the imposition of the European Medicines Agency to include safety information on immune thrombocytopenia.

Vaccinated individuals should seek immediate medical assistance if they experience the following symptoms after vaccination:⁹

- unexplained bleeding
- unexplained bruising
- small purplish spots beyond the site of vaccination
- shortness of breath
- chest pain
- leg pain and/or swelling
- persistent abdominal pain

Four (4) cases of immune thrombocytopenia have been reported. Two (2) cases have been assessed as coincidental or not related to the vaccine and two (2) cases are currently being reviewed. The FDA together with the Epidemiology Bureau of the DOH shall continue to monitor vaccine safety ensuring that the benefits always outweigh the risks.

⁹ Health Canada is updating the labels of the Janssen and Vaxzevria (AstraZeneca) COVID-19 vaccines
<https://recalls-rappels.canada.ca/en/alert-recall/health-canada-updating-labels-janssen-and-vaxzevria-astrazeneca-covid-19-vaccines>

Cases of hospitalization

One of the criteria for serious adverse reaction is hospitalization or extended hospital stay. Reports of adverse reaction that results in hospitalization does not necessarily mean that vaccine caused the reaction. An Expert Committee reviews and assesses whether the vaccine caused the reaction. Based on the reports received, the hospitalization-reporting rate is 2.83 per 100,000 doses administered. Commonly reported causes of hospitalization include pyrexia, cough, dyspnea, and headache.

Reports involving death

As of 27 February 2022, 1,918 fatal events were received. Reports of fatal events does not necessarily mean that the vaccine caused the events. Underlying conditions or pre-existing

medical conditions causing fatal events are usually coincidental on the use of the vaccine. It is expected that reports of fatal events will rise as the vaccination program covers more people including those with undiagnosed illness, underlying comorbidities, and pre-existing medical conditions.

The vaccinees reported to have fatal events were aged 12 years and above. The mean age of the fatal cases was 61.17 years. 62.10% (1,191) of the fatal cases were from age group 60 years and above, 24.61% (472) from age group of 40-59 years, 11.00% (211) from age group 18-39 years, 1.04% (20) from age group 12-17 years of age and 1.25% (24) were not identified to what age group they are classified.

Most of these events occurred in persons with multiple existing comorbidities. These include cardiovascular diseases, ischemic heart diseases, cerebrovascular diseases, cancer, diabetes, and infections including pneumonia. There were cases of confirmed COVID-19 infections leading to severe cases with fatal outcomes.

Number of suspected adverse reactions per category

A total of 92,215 case reports consisting of 198,892 suspected adverse reactions were received from the start of the vaccination program. More than one suspected adverse reaction might be reported in a single case. Suspected adverse reactions were coded using the Medical Dictionary for Regulatory Activities (MedDRA) terminology to allow international comparison of reports.

The data presented below are categorized by System Organ Class (SOC), the highest in the hierarchy of MedDRA. They are grouped by manifestation site (e.g. gastrointestinal, cardiac) and etiology (e.g. infections, examinations).

Reactions to inactivated vaccines

- CoronaVac
- COVID-19 Vaccine Sinopharm

Classification	Number of suspected reactions
General symptoms & reactions in the administration site <i>E.g. Pain and reaction in the injection site, chills, discomfort, fever, fatigue</i>	14,185
Cardiac symptoms <i>E.g. Palpitations, bradycardia, tachycardia</i>	950
Ear symptoms <i>E.g. Ear swelling, vertigo, tinnitus, ear discomfort</i>	61
Endocrine symptoms <i>E.g. Adrenal insufficiency, goiter, thyroid symptoms</i>	5
Examinations <i>E.g. Increased blood pressure, increased heart rate, blood glucose increased, SARS-CoV-2 test</i>	12,556

Eye symptoms <i>E.g. Eye itchiness, blurred vision, eye pain, eye swelling</i>	281
Gastrointestinal symptoms <i>E.g. Abdominal pain, diarrhea, nausea, vomiting, dry mouth, lip swelling</i>	3,741
Hepatobiliary symptoms <i>E.g. Jaundice</i>	10
Immune system symptoms <i>E.g. Allergic reactions, hypersensitivity</i>	202
Infections <i>E.g. Cold symptoms, rhinitis</i>	4,047
Metabolism and nutrition-related symptoms <i>E.g. Decreased appetite, increased appetite, starvation, dehydration</i>	486
Musculoskeletal symptoms <i>E.g. Back pain, joint pain, pain in extremities, muscle pain, muscle spasms</i>	2,305
Neoplasm <i>E.g. Liver cancer, endometrial cancer, uterine myoma</i>	10
Neurological symptoms <i>E.g. Dizziness, headache, syncope</i>	9,161
Pregnancy, puerperium, and perinatal conditions <i>E.g. Abortion, hemorrhage</i>	26
Procedural symptoms <i>E.g. Procedural hypertension, vaccination adverse reaction</i>	128
Psychiatric symptoms <i>E.g. Feeling anxious, insomnia, nervousness, disorientation</i>	173
Renal and urinary symptoms <i>E.g. Urine coloring yellow, urine frequency</i>	60
Reproductive symptoms <i>E.g. Vaginal bleeding, vaginal spotting</i>	67
Respiratory symptoms <i>E.g. Cough, nasal congestion, throat irritation</i>	7,045
Skin symptoms <i>E.g. Cold sweat, rash, redness</i>	4,406
Social circumstances <i>E.g. Hearing disability, walking disability</i>	3
Surgical and medical procedures <i>E.g. Tumor debulking, nasolabial flap</i>	12
Symptoms in blood and lymphatic system <i>E.g. Pain in the lymph nodes</i>	47
Vascular symptoms <i>E.g. Flashes, low blood pressure</i>	385




Figure 34. Suspected adverse reaction distribution by SOC for inactivated vaccine

As shown in Figure 3, the SOC which consists of the greatest number of reports were examinations (12,436), followed by general symptoms and reactions in the administration site (10,498), neurological symptoms (7,676), respiratory symptoms (5,370), infections (3,495), skin symptoms (3,318), gastrointestinal symptoms (2,803), musculoskeletal symptoms (2,000), cardiac symptoms (913), and metabolism and nutrition-related symptoms (482).

The top reported events for CoronaVac are:

- blood pressure increased (36.52%)
- pyrexia (13.81%)
- headache (12.70%)
- cough (10.50%)
- vaccination/injection site pain (10.21%)
- dizziness (6.71%)
- rash (6.66%)
- nasopharyngitis (5.89%)
- malaise (4.95%)
- dyspnea (4.81%)

The top reported events for COVID-19 Vaccine Sinopharm are:

- pyrexia (23.70%)
- cough (15.64%)

- dizziness (14.22%)
- headache (10.90%)
- blood pressure increased (8.53%)
- rash (8.06%)
- nasopharyngitis (7.11%)
- dyspnea (6.16%)
- fatigue (5.69%)
- COVID-19 (4.74%), pruritus (4.74%)

Reactions to non-replicating viral vector vaccines

- COVID-19 vaccine AstraZeneca
- Sputnik V
- Janssen COVID-19 Vaccine

Classification	Number of suspected reactions
General symptoms & reactions in the administration site <i>E.g. Pain and reaction in the injection site, chills, discomfort, fever, fatigue</i>	46,028
Cardiac symptoms <i>E.g. Palpitations, bradycardia, tachycardia</i>	816
Congenital, familial, and genetic disorder <i>E.g. Polycystic kidney</i>	3
Ear symptoms <i>E.g. Ear swelling, vertigo, tinnitus, ear discomfort</i>	71
Endocrine symptoms <i>E.g. Adrenal insufficiency, goiter, thyroid symptoms</i>	3
Examinations <i>E.g. Increased blood pressure, increased heart rate, blood glucose increased, SARS-CoV-2 test</i>	7,295
Eye symptoms <i>E.g. Eye itchiness, blurred vision, eye pain, eye swelling</i>	521
Gastrointestinal symptoms <i>E.g. Abdominal pain, diarrhea, nausea, vomiting, lip swelling</i>	5,204
Hepatobiliary symptoms <i>E.g. Jaundice</i>	14
Immune system symptoms <i>E.g. Allergic reactions, hypersensitivity</i>	286
Infections <i>E.g. Cold symptoms, rhinitis</i>	3,555
Metabolism and nutrition-related symptoms <i>E.g. Decreased appetite, increased appetite, starvation, dehydration</i>	790
Musculoskeletal symptoms <i>E.g. Back pain, joint pain, pain in extremities, muscle pain, muscle spasms</i>	9,289
Neoplasms <i>E.g. Liver cancer, endometrial cancer, uterine myoma</i>	6
Neurological symptoms <i>E.g. Dizziness, headache, syncope</i>	17,418
Pregnancy, puerperium, and perinatal conditions <i>E.g. Abortion, hemorrhage</i>	28

Procedural symptoms <i>E.g. Procedural hypertension, vaccination adverse reaction</i>	207
Psychiatric symptoms <i>E.g. Feeling anxious, insomnia, nervousness, disorientation</i>	122
Renal and urinary symptoms <i>E.g. Urine coloring yellow, urine frequency</i>	67
Reproductive symptoms <i>E.g. Vaginal bleeding, vaginal spotting</i>	74
Respiratory symptoms <i>E.g. Cough, nasal congestion, throat irritation</i>	5,311
Skin symptoms <i>E.g. Cold sweat, rash, redness</i>	3,828
Social circumstances <i>E.g. Hearing disability, walking disability</i>	2
Surgical and medical procedures <i>E.g. Tumor debulking, nasolabial flap</i>	12
Symptoms in blood and lymphatic system <i>E.g. Pain in the lymph nodes</i>	80
Vascular symptoms <i>E.g. Flushes, low blood pressure</i>	382




Figure 4. Suspected adverse reaction distribution by SOC for viral vector vaccines

As shown in Figure 4, the SOC which consists of the greatest number of reports were general symptoms and reactions in the administration site (24,437), followed by neurological symptoms (15,119), musculoskeletal symptoms (7,540), examinations (7,248), respiratory symptoms (4,232), gastrointestinal symptoms (3,998), infections (3,082), skin symptoms (2,947), metabolism and nutrition-related symptom (780), and cardiac symptoms (778).

The top reported events for COVID-19 Vaccine AstraZeneca are:

- pyrexia (39.39%)
- headache (32.68%)
- vaccination/injection site pain (22.49%)
- malaise (20.55%)
- blood pressure increased (15.79%)
- chills (15.74%)
- myalgia (14.95%)
- fatigue (11.19%)
- arthralgia (7.39%)
- dizziness (6.45%)

The top reported events for Sputnik V are:

- blood pressure increased (52.70%)
- pyrexia (13.14%)
- headache (8.48%)
- cough (7.00%)
- dizziness (5.41%), nasopharyngitis (5.41%), rash (5.41%)
- vaccination/injection site pain (5.16%)
- COVID-19 (4.30%), heart rate increased (4.30%)
- chills (3.81%)
- dyspnea (3.07%)
- nausea (2.83%)

The top reported events for Janssen COVID-19 Vaccine are:

- blood pressure increased (25.18%)
- pyrexia (19.60%)
- vaccination/injection site pain (15.43%)
- cough (11.15%)
- headache (11.03%)
- dyspnea (6.49%)
- dizziness (6.23%)
- hypoesthesia (5.35%)
- malaise (5.18%)
- COVID-19 (5.16%)

Reactions to mRNA vaccines

- Comirnaty
- COVID-19 Vaccine Moderna

Classification	Number of suspected reactions
General symptoms & reactions in the administration site <i>E.g. Pain and reaction in the injection site, chills, discomfort, fever, fatigue</i>	12,737
Cardiac symptoms <i>E.g. Palpitations, bradycardia, tachycardia</i>	666
Congenital, familial, and genetic disorder <i>E.g. Skull malformation</i>	3
Ear symptoms <i>E.g. Ear swelling, vertigo, tinnitus, ear discomfort</i>	39
Endocrine symptoms <i>E.g. Adrenal insufficiency, goiter, thyroid symptoms</i>	3
Examinations <i>E.g. Increased blood pressure, increased heart rate, blood glucose increased, SARS-CoV-2 test</i>	4,625
Eye symptoms <i>E.g. Eye itchiness, blurred vision, eye pain, eye swelling</i>	222
Gastrointestinal symptoms <i>E.g. Abdominal pain, diarrhea, nausea, vomiting, lip swelling</i>	2,246
Hepatobiliary symptoms <i>E.g. Jaundice</i>	8
Immune system symptoms <i>E.g. Allergic reactions, hypersensitivity</i>	144
Infections <i>E.g. Cold symptoms, rhinitis</i>	1,667
Metabolism and nutrition-related symptoms <i>E.g. Decreased appetite, increased appetite, starvation, dehydration</i>	232
Musculoskeletal symptoms <i>E.g. Back pain, joint pain, pain in extremities, muscle pain, muscle spasms</i>	1,873
Neoplasms <i>E.g. Liver cancer, endometrial cancer, uterine myoma</i>	6
Neurological symptoms <i>E.g. Dizziness, headache, syncope</i>	5,958
Pregnancy, puerperium, and perinatal conditions <i>E.g. Abortion, hemorrhage</i>	67
Procedural symptoms <i>E.g. Procedural hypertension, vaccination adverse reaction</i>	152
Psychiatric symptoms <i>E.g. Feeling anxious, insomnia, nervousness, disorientation</i>	121
Renal and urinary symptoms <i>E.g. Urine coloring yellow, urine frequency</i>	35
Reproductive symptoms <i>E.g. Vaginal bleeding, vaginal spotting</i>	78
Respiratory symptoms <i>E.g. Cough, nasal congestion, throat irritation</i>	3,146

Skin symptoms <i>E.g. Cold sweat, rash, redness</i>	2,712
Social circumstances <i>E.g. Hearing disability, walking disability</i>	6
Surgical and medical procedure <i>E.g. Chemotherapy</i>	54
Symptoms in blood and lymphatic system <i>E.g. Pain in the lymph nodes</i>	98
Vascular symptoms <i>E.g. Flushes, low blood pressure</i>	200




Figure 5. Suspected adverse reaction distribution by SOC for mRNA vaccine

As shown in Figure 5, the SOC which consists of the greatest number of reports were general symptoms and reactions in the administration site (8,105), followed by neurological symptoms (5,000), examinations (4,565), respiratory symptoms (2,445), skin symptoms (2,098), gastrointestinal symptoms (1,662), musculoskeletal symptoms (1,566), infections (1,458), cardiac symptoms (652), and metabolism and nutrition-related symptom (230).

The top reported events for Comirnaty are:

- blood pressure increased (23.41%)
- pyrexia (18.54%)
- vaccination/injection site pain (17.12%)

- headache (13.63%)
- dizziness (10.48%)
- rash (7.62%)
- cough (7.51%)
- malaise (6.14%)
- nasopharyngitis (4.82%)
- dyspnea (4.60%)

The top reported events for COVID-19 vaccine Moderna are:

- pyrexia (30.16%)
- vaccination/injection site pain (20.67%)
- blood pressure increased (20.53%)
- headache (16.23%)
- cough (8.22%)
- chills (8.05%)
- dizziness (7.55%)
- rash (7.02%)
- malaise (6.06%)
- myalgia (5.77%)

Outcome of suspected adverse reactions

The outcome of cases of suspected adverse reactions to COVID-19 vaccines is shown in Figure 6. Overall, most of the reported cases have *recovered/resolved* (75.15%), although there were few cases who have *recovered but with sequelae* (0.04%). Over 13% of the cases are *recovering/resolving* while more than 1% have *not recovered/not resolved* at the time of reporting. A proportion of 2.08% were reported with fatal outcomes as discussed in the section reports involving death.




Figure 6. Case outcome

Reporting of suspected adverse reactions following vaccination

Individuals who have received their COVID-19 vaccination shots can report any suspected adverse reaction to any of the following:

- Immunization site where you were vaccinated
- Directly to the vaccine manufacturer or emergency use authorization holder
 - [Sinovac – CoronaVac & Covaxin](#)
 - [AstraZeneca – COVID-19 Vaccine AstraZeneca](#)
 - [Gamaleya – Sputnik V & Sputnik Light](#)
 - [Pfizer – Comirnaty](#)
 - [Zuellig – COVID-19 Vaccine Moderna](#)
 - Johnson & Johnson – Janssen COVID-19 Vaccine
 - Faberco – Covovax
- [FDA online reporting system](#)

Kindly **report only to one** of the above to avoid duplication of reports.