

FACT SHEET FOR RECIPIENTS AND CAREGIVERS

EMERGENCY USE AUTHORIZATION (EUA) OF THE JANSSEN COVID-19 VACCINE TO PREVENT CORONAVIRUS DISEASE 2019 (COVID-19) IN INDIVIDUALS 18 YEARS OF AGE AND OLDER

You are being offered the Janssen COVID-19 Vaccine to prevent Coronavirus Disease 2019 (COVID-19) caused by SARS-CoV-2. This Fact Sheet contains information to help you understand the risks and benefits of receiving the Janssen COVID-19 Vaccine, which you may receive because there is currently a pandemic of COVID-19.

The Janssen COVID-19 Vaccine may prevent you from getting COVID-19. There is no Philippine Food and Drug Administration (PFDA) approved vaccine to prevent COVID-19.

Read this Fact Sheet for information about the Janssen COVID-19 Vaccine. Talk to the vaccination provider if you have questions. It is your choice to receive the Janssen COVID-19 Vaccine.

The Janssen COVID-19 Vaccine has received EUA from PFDA to provide:

- A single dose primary vaccination to individuals 18 years of age and older.
- A single booster dose to individuals 18 years of age and older who have completed a primary vaccination with the Janssen COVID-19 Vaccine.
- A single booster dose to individuals 18 years of age and older who have completed primary vaccination with a different authorized COVID-19 vaccine.

The Janssen COVID-19 Vaccine may not protect everyone.

This Fact Sheet may have been updated. For the most recent Fact Sheet, please visit <https://vaxcheck.jnj/> or <https://www.janssencovid19vaccine.com.ph/>.

WHAT YOU NEED TO KNOW BEFORE YOU GET THIS VACCINE

WHAT IS COVID-19?

COVID-19 is caused by a coronavirus called SARS-CoV-2. This type of coronavirus has not been seen before. You can get COVID-19 through contact with another person who has the virus. It is predominantly a respiratory illness that can affect other organs. People with COVID-19 have had a wide range of symptoms reported, ranging from mild symptoms to severe illness. Symptoms may appear 2 to 14 days after exposure to the virus. Common symptoms may include: fever or chills; cough; shortness of breath; fatigue; muscle or body aches; headache; new loss of taste or smell; sore throat; congestion or runny nose; nausea or vomiting; diarrhea.

WHAT IS THE JANSSEN COVID-19 VACCINE?

The Janssen COVID-19 Vaccine is an unapproved vaccine that may prevent COVID-19. There is no PFDA-approved vaccine to prevent COVID-19.

The PFDA has authorized the emergency use of the Janssen COVID-19 Vaccine to prevent COVID-19 in individuals 18 years of age and older under an Emergency Use Authorization (EUA).

For more information on EUA, see the “**What is an Emergency Use Authorization (EUA)?**” section at the end of this Fact Sheet.

WHAT SHOULD YOU MENTION TO YOUR VACCINATION PROVIDER BEFORE YOU GET THE JANSSEN COVID-19 VACCINE?

Tell the vaccination provider about all of your medical conditions, including if you:

- have any allergies,
- have a fever,
- have a bleeding disorder or are on a blood thinner,
- have ever had a low level of platelets (blood cells that help your body stop bleeding),
- are immunocompromised or are on a medicine that affects your immune system,
- are pregnant or plan to become pregnant,
- are breastfeeding,
- have received another COVID-19 vaccine,
- have ever fainted in association with an injection

WHO SHOULD GET THE JANSSEN COVID-19 VACCINE?

PFDA has authorized the emergency use of the Janssen COVID-19 Vaccine in individuals 18 years of age and older.

WHO SHOULD NOT GET THE JANSSEN COVID-19 VACCINE?

You should not get the Janssen COVID-19 Vaccine if you:

- had a severe allergic reaction after a previous dose of this vaccine.
- had a severe allergic reaction to any ingredient of this vaccine.
- had a blood clot along with a low level of platelets (blood cells that help your body stop bleeding) following Janssen COVID-19 Vaccine or following any other adenovirus-vectored COVID-19 vaccine

WHAT ARE THE INGREDIENTS IN THE JANSSEN COVID-19 VACCINE?

The Janssen COVID-19 Vaccine includes the following ingredients: recombinant, replication-incompetent adenovirus type 26 expressing the SARS-CoV-2 spike protein, citric acid monohydrate, trisodium citrate dihydrate, ethanol, 2-hydroxypropyl- β -cyclodextrin (HBCD), polysorbate-80, sodium chloride.

HOW IS THE JANSSEN COVID -19 VACCINE GIVEN?

The Janssen COVID-19 Vaccine will be given to you as an injection into the muscle.

Primary Vaccination

The Janssen COVID-19 Vaccine is administered as a **single dose**.

Booster Dose

- A single booster dose of the Janssen COVID-19 Vaccine may be administered at least two months after primary vaccination with the Janssen COVID-19 Vaccine.
- A single booster dose of the Janssen COVID-19 Vaccine may be administered to individuals 18 years of age and older who have completed primary vaccination with a different authorized COVID-19 vaccine. Please check with your health care provider regarding timing of the booster dose.

HAS THE JANSSEN COVID-19 VACCINE BEEN USED BEFORE?

The Janssen COVID-19 Vaccine is an unapproved vaccine. In an ongoing clinical trials, more than 61,000 individuals 18 years of age and older have received the Janssen COVID-19 Vaccine. Millions of individuals have received the vaccine under EUA since February 27, 2021.

WHAT ARE THE BENEFITS OF THE JANSSEN COVID-19 VACCINE?

The Janssen COVID-19 Vaccine has been shown to prevent COVID-19. The duration of protection against COVID-19 is currently unknown.

WHAT ARE THE RISKS OF THE JANSSEN COVID-19 VACCINE?

Side effects that have been reported with the Janssen COVID-19 Vaccine include:

- Injection site reactions: pain, redness of the skin and swelling.
- General side effects: headache, feeling very tired, muscle aches, nausea, and fever.
- Swollen lymph nodes
- Blood clots
- Unusual feeling in the skin (such as tingling or a crawling feeling) (paresthesia), decreased feeling or sensitivity, especially in the skin (hypoesthesia).
- Persistent ringing in the ears (tinnitus)
- Diarrhea, vomiting.

Severe Allergic Reactions

There is a remote chance that the Janssen COVID-19 Vaccine could cause a severe allergic reaction. A severe allergic reaction would usually occur within a few minutes to one hour after

getting a dose of the Janssen COVID-19 Vaccine. For this reason, your vaccination provider may ask you to stay at the place where you received your vaccine for monitoring after vaccination. Signs of a severe allergic reaction can include:

- Difficulty breathing,
- Swelling of your face and throat,
- A fast heartbeat,
- A bad rash all over your body,
- Dizziness and weakness.

Blood Clots with Low Levels of Platelets

Blood clots involving blood vessels in the brain, lungs, abdomen, and legs along with low levels of platelets (blood cells that help your body stop bleeding), have occurred in some people who have received the Janssen COVID-19 Vaccine. In people who developed these blood clots and low levels of platelets, symptoms began approximately one to two-weeks after vaccination. Blood clots with low levels of platelets following the Janssen COVID-19 Vaccine have been reported in males and females, across a wide age range of individuals 18 years and older; reporting has been highest in females ages 30 through 49 years (about 1 case for every 100,000 vaccine doses administered), and about 1 out of every 7 cases has been fatal. You should seek medical attention right away if you have any of the following symptoms after receiving the Janssen COVID-19 Vaccine:

- Shortness of breath,
- Chest pain,
- Leg swelling,
- Persistent abdominal pain,
- Severe or persistent headaches or blurred vision,
- Easy bruising or tiny blood spots under the skin beyond the site of the injection.

Immune Thrombocytopenia (ITP)

Immune Thrombocytopenia (ITP) is a disorder that can cause easy or excessive bruising and bleeding due to very low levels of platelets. ITP has occurred in some people who have received the Janssen COVID-19 Vaccine. In most of these people, symptoms began within 42 days following receipt of the Janssen COVID-19 Vaccine. The chance of having this occur is very low. If you have ever had a diagnosis of ITP, talk to your vaccination provider before you get the Janssen COVID-19 Vaccine. You should seek medical attention right away if you develop any of the following symptoms after receiving the Janssen COVID-19 Vaccine:

- Easy or excessive bruising or tiny blood spots under the skin beyond the site of the injection,
- Unusual or excessive bleeding.

Guillain Barré Syndrome

Guillain Barré syndrome (a neurological disorder in which the body's immune system damages nerve cells, causing muscle weakness and sometimes paralysis) has occurred in some people who have received the Janssen COVID-19 Vaccine. In most of these people, symptoms began within 42 days following receipt of the Janssen COVID-19 Vaccine. The chance of having this occur is very low. You should seek medical attention right away if you develop any of the following symptoms after receiving the Janssen COVID-19 Vaccine:

- Weakness or tingling sensations, especially in the legs or arms, that's worsening and spreading to other parts of the body.
- Difficulty walking.
- Difficulty with facial movements, including speaking, chewing or swallowing.
- Double vision or inability to move eyes.
- Difficulty with bladder control or bowel function.

These may not be all the possible side effects of the Janssen COVID-19 Vaccine. Serious and unexpected effects may occur. The Janssen COVID-19 Vaccine is still being studied in clinical trials.

WHAT SHOULD I DO ABOUT SIDE EFFECTS?

If you experience a severe allergic reaction, go to the nearest hospital.

Call the vaccination provider or your healthcare provider if you have any side effects that bother you or do not go away.

Report vaccine side effects to **PFDA** via Telephone. no: +632 8 857-1900 or report online to <https://www.fda.gov.ph/pharmacovigilance/>.

WHAT IF I DECIDE NOT TO GET THE JANSSEN COVID-19 VACCINE?

It is your choice to receive or not receive the Janssen COVID-19 Vaccine. Should you decide not to receive it, it will not change your standard medical care.

ARE OTHER CHOICES AVAILABLE FOR PREVENTING COVID-19 BESIDES JANSSEN COVID-19 VACCINE?

Currently, there is no PFDA approved alternative vaccine available for prevention of COVID-19. Other vaccines to prevent COVID-19 may be available under Emergency Use Authorization.

CAN I RECEIVE THE JANSSEN COVID-19 VACCINE WITH OTHER VACCINES?

There is no information on the use of the Janssen COVID-19 Vaccine with other vaccines.

WHAT IF I AM PREGNANT OR BREASTFEEDING?

If you are pregnant or breastfeeding, discuss your options with your healthcare provider.

WILL THE JANSSEN COVID-19 VACCINE GIVE ME COVID-19?

No. The Janssen COVID-19 Vaccine does not contain SARS-CoV-2 and cannot give you COVID-19.

KEEP YOUR VACCINATION CARD

When you receive the Janssen COVID-19 Vaccine, please ask a vaccination card to document the name of the vaccine and date of when you received the vaccine. Please keep this record.

ADDITIONAL INFORMATION

If you have questions or to access the most recent Janssen COVID-19 Vaccine Fact Sheets, visit the websites <https://vaxcheck.jnj/> or <https://www.janssencovid19vaccine.com.ph/> or call the telephone numbers provided below:

- Toll-free number (PLDT): 1800-1110-2906
- Toll-free number (Globe): 1800-8909-4047

HOW CAN I LEARN MORE?

- Ask the vaccination provider.
- Visit PFDA at <https://www.fda.gov.ph/fda-circular-no-2020-036-guidelines-on-the-issuance-of-emergency-use-authorization-for-drugs-and-vaccines-for-covid-19/>

Contact your local or state public health department.

WHAT IS AN EMERGENCY USE AUTHORIZATION (EUA)?

The PFDA has made the Janssen COVID-19 Vaccine available under an emergency access mechanism called an EUA. The EUA is supported by a Department of Health declaration that circumstances exist to justify the emergency use of drugs and biological products during the COVID-19 pandemic.

For the definition of EUA please refer to <https://www.fda.gov.ph/fda-circular-no-2020-036-guidelines-on-the-issuance-of-emergency-use-authorization-for-drugs-and-vaccines-for-covid-19/>

The Janssen COVID-19 Vaccine has not undergone the same type of review as an FDA-approved or cleared product. FDA may issue an EUA when certain criteria are met, which includes that there are no adequate, approved, and available alternatives. In addition, the FDA decision is based on the totality of scientific evidence available showing that the product may be effective to prevent COVID-19 during the COVID-19 pandemic and that the known and potential benefits of the

product outweigh the known and potential risks of the product. All of these criteria must be met to allow for the product to be used during the COVID-19 pandemic.

The EUA for the Janssen COVID-19 Vaccine is in effect for the duration of the COVID-19 declaration justifying emergency use of these products, unless terminated or revoked (after which the products may no longer be used).



© 2021 Janssen Pharmaceutical Companies

For more information, go to <https://vaxcheck.jnj/>.

Revised: Apr/6/2022 (based on US Fact Sheet Jan/11/2022)