



Reports of Suspected Adverse Reaction to COVID-19 Vaccines (01 March to 05 December 2021)

Contents

About the report	2
Summary	3
COVID-19 vaccines with Emergency Use Authorization in the Philippines	4
Statistics regarding reports of suspected adverse reactions	4
Distribution of reports of adverse reaction for each vaccine.....	5
Demographics	5
Vaccination in adolescent population	6
Booster shots	7
National Vaccination Day	7
Reports of suspected serious adverse reaction	7
Number of suspected adverse reactions per category	12
Reactions to inactivated vaccines	12
Reactions to non-replicating viral vector vaccines	15
Reactions to mRNA vaccines	18
Outcome of suspected adverse reaction	20
Reporting of suspected adverse reactions following vaccination	20



About the report

- A summary is presented below of all received suspected adverse reaction reports following COVID-19 vaccination from 01 March 2021, the date when the first vaccine became available, up to 05 December 2021.
- Seven (7) vaccines under Emergency Use Authorization (EUA) are currently being used in the vaccination program: CoronaVac, COVID-19 Vaccine AstraZeneca, Sputnik V, Comirnaty, COVID-19 Vaccine Moderna, Janssen COVID-19 Vaccine, and COVID-19 Vaccine Sinopharm.
- Data are based on the VigiFlow, the national database of adverse reactions in the Philippines. It includes reports from various epidemiology surveillance units (ESUs) of the Department of Health (DOH), vaccination sites, hospitals, patients/consumers, and EUA holders.
- Symptoms or diseases that occur after vaccination are reported if there is a *suspicion* of a possible link. However, it cannot be assumed that there is a causal relationship between the suspected adverse reaction and the vaccine.
- This report contains all suspected adverse reactions regardless of any possible causal relationship.
- Additional information may become available in individual case reports at any time, which may change the assessment and figures presented.
- Adverse reaction reports are necessary for the safety assessment of the vaccines, making sure that the benefits always outweigh the risks.
- Reports are constantly reviewed and monitored for the possible emergence/identification of unknown adverse reactions also known as signal. If a signal is identified, investigations, regulatory actions, and timely communication is performed by the FDA and DOH.
- A weekly report is published to summarize reported adverse reactions to the COVID-19 vaccines.

Summary

This report is based on an assessment of adverse reaction reports received by 05 December 2021. As per benefit-risk assessment, these reports do not provide a basis for revising the current recommendations regarding the use of COVID-19 vaccines.

A report of adverse reaction does not necessarily mean that the vaccine caused the reactions. A mere suspicion may also be reported. Undiagnosed illness, underlying comorbidities, and pre-existing medical conditions unrelated to vaccination can be factors in reporting adverse reactions. The relative numbers should not be used to compare the safety of different vaccines.

Like any other vaccines, COVID-19 vaccines may cause adverse reactions in some people while others may not experience any adverse reaction. It is possible for several persons to experience the same adverse event but for the report to be serious for one person and non-serious for another person. Most of the reported reactions are generally in line with what is described in the product information and labels. Such reports are minor adverse reactions, which include body pain, chills, fatigue, fever, headache, nausea, and pain in the injection site. These usually appear on the first or second day of vaccination and may last for 2-3 days. Most people tolerate these adverse reactions while others experience greater discomfort.

Serious adverse reactions have also been reported. The FDA together with other public health partners are continuously monitoring the adverse experience as more people are being vaccinated with COVID-19 vaccines. Such monitoring will provide reassurance that the vaccines are safe and effective for use.

Considering the post-authorization experience on the use of COVID-19 Vaccine AstraZeneca and Janssen COVID-19 Vaccine of other countries, information on the very rare and serious adverse events of thrombosis and thrombocytopenia and in some cases accompanied by bleeding have been revised under the special warning and precautions for use in their product labels. In addition, capillary leak syndrome and Guillain-Barré syndrome were added as a rare new side effect.

The label of mRNA vaccines Moderna and Comirnaty (Pfizer-BioNTech) have been revised to include imposition of the European Medicines Agency and the USFDA to include safety information on myocarditis and pericarditis.

The immunization program expanded its coverage to include adolescent individuals (12 to 17 years old). Pfizer and Moderna are the only vaccines with EUA for the said population. The roll-out started on the second week of October. Likewise, monitoring of suspected adverse reactions has extended to include such population.

Seven (7) vaccines are currently used in the immunization program. These include CoronaVac, COVID-19 Vaccine AstraZeneca, Sputnik V, Comirnaty, COVID-19 Vaccine Moderna, Janssen COVID-19 Vaccine, and COVID-19 Vaccine Sinopharm. Supplies of vaccines are either procured by the government and/or private sector or supplied under the COVAX facility.

COVID-19 vaccines with Emergency Use Authorization in the Philippines

At present, the FDA granted [nine \(9\) COVID-19 vaccines with emergency use authorization](#):

- Pfizer-BioNTech COVID-19 Vaccine (Comirnaty)
- ChAdOx1-S [recombinant] (COVID-19 Vaccine AstraZeneca)
- SARS-CoV-2 Vaccine (Vero Cell), Inactivated (CoronaVac)
- Gam-COVID-Vac (Sputnik V & Sputnik Light)
- Ad26.COV2-S [recombinant] (Janssen COVID-19 Vaccine)
- Whole Virion Inactivated Corona Virus (Covaxin)
- COVID-19 mRNA Vaccine [nucleoside modified] (COVID-19 Vaccine Moderna)
- Inactivated COVID-19 Vaccine (Vero Cell) (COVID-19 Vaccine Sinopharm BIBP/Wuhan)
- SARS-CoV-2 rS Protein Nanoparticle Vaccine [Covovax]

Various vaccine platforms have been approved for use in the Philippines. Comirnaty and COVID-19 Vaccine Moderna are mRNA vaccines; COVID-19 Vaccine AstraZeneca and Janssen COVID-19 Vaccine are non-replicating viral vector vaccines while Sputnik V uses the same technology having two (2) different (dose) components of viral vectors and Spuntik Light having single dose component; CoronaVac, Covaxin, and COVID-19 Vaccine Sinopharm BIBP/Wuhan are inactivated vaccines; and Covovax is a protein subunit vaccine. All are administered in two doses within an interval of a few weeks except for Janssen COVID-19 Vaccine and Sputnik Light, which are administered as a single-dose.

Statistics regarding reports of suspected adverse reactions

As of 05 December 2021, more than 91.7 million doses of COVID-19 vaccines (either CoronaVac, COVID-19 Vaccine AstraZeneca, Sputnik V, Comirnaty, COVID-19 Vaccine Moderna, Janssen COVID-19 Vaccine, or COVID-19 Vaccine Sinopharm) were already administered. Over 38.1 million individuals are now fully vaccinated (either given a single-dose or 2-dose vaccine course) while more than 18.5 million are partly vaccinated waiting for their second dose to be administered. A total of 509,944 individuals already received their third or additional doses (either homologous or heterologous booster doses). A total of 80,644 suspected adverse reaction reports were received, evaluated, and analyzed by the FDA. To disaggregate, 30,737 have been reported for CoronaVac, 33,285 for COVID-19 Vaccine AstraZeneca, 767 for Sputnik V, 8,839 for Comirnaty, 3,248 for COVID-19 Vaccine Moderna, 3,614 for Janssen COVID-19 Vaccine, and 154 for COVID-19 Vaccine Sinopharm.

Table 1. Data on vaccination and suspected adverse reaction reports.

Indicators	Value
Total number of doses administered	91,777,433
No. of fully vaccinated individuals	38,168,603
No. of individuals partly vaccinated	18,531,877
No. of individuals with booster shots	509,944

No. of suspected adverse reaction reports	80,644 (0.09% of doses administered)
No. of suspected serious adverse reaction reports	4,077 (0.004% of doses administered)

Distribution of reports of adverse reactions for each vaccine

Data shown below are cumulative reports from the start of the vaccination program on 01 March 2021 up until 05 December 2021.

Table 2. Distribution of reports of adverse reactions for each vaccine

Vaccine	Date started	Total vaccine doses administered ^b	Number of fully vaccinated individuals ^b	Number of individuals partly vaccinated	Number of individuals with booster shots	Total number of reports ^a	Reports of non-serious events	Reports of serious events
CoronaVac	01 Mar 2021	37,731,431	16,717,306	4,269,826	26,993	30,737	28,839	1,898
AstraZeneca	07 Mar 2021	12,783,998	4,649,668	3,426,942	57,720	33,285	32,248	1,037
Sputnik V	04 May 2021	779,686	301,728	176,084	146	767	740	27
Comirnaty	13 May 2021	26,172,932	8,867,328	8,077,998	360,278	8,839	8,406	433
Moderna	30 June 2021	9,748,111	3,587,206	2,508,948	64,751	3,248	3,044	204
Janssen	20 July 2021	3,601,594	3,601,594	-	-	3,614	3,150	464
Sinopharm	25 Aug 2021	959,681	443,773	72,079	56	154	140	14
TOTAL		91,777,433	38,168,603	18,531,877	509,944	80,644	76,567	4,077

Data source: ^aVigiFlow, ^bNVOC daily report as 6PM, 05 December 2021

Notes: Additional information may become available in individual cases, which may change the figures presented

^bAn individual is considered partly vaccinated if they have received only one dose of a two-dose vaccine course. An individual is considered fully vaccinated if they have received a single-dose vaccine or both doses of a two-dose vaccine

^cData concerning various vaccines are not directly comparable. COVID-19 vaccines profile varies, they have not been used for equal periods of time and they have been administered to number of people with different profiles including various age and sex.

Demographics

The figures below provide a descriptive overview of the population who have experienced adverse reactions to COVID-19 vaccines. Figure 1 and Figure 2 shows the distribution of reports by gender and age. Click [here](#) to show disaggregated data.




Figure 1. Report distribution by gender




Figure 2. Report distribution by age

The early phase of the vaccination program is intended for the frontline health workers, thus, the high number in the female and younger population might be associated with the data that our health system is dominated by female (75%) and young adults under the age of 35 (65%).¹ An increasing number of reports from the age group 40 years and above have been observed in the past few weeks of the vaccination program. This may be attributed to the coverage of priority groups of senior citizens and individuals with comorbidities.

Relative to the inclusion of the frontline personnel in the priority groups, the observed increasing number of reports in the male population may be attributed to the vaccine coverage and statistics that more males are employed than females (6 in every 10).²

¹ Human Resource for Health in the Time of the COVID-19 Pandemic: Does the Philippines Have Enough?

<https://www.drdf.org.ph/sites/default/files/pdf/COVID-19-Research-Brief-08.pdf>

² Employment situation in July 2018, Philippine Statistics Authority

<https://psa.gov.ph/statistics/survey/labor-and-employment/labor-force-survey/title/Employment%20Situation%20in%20July%202018>

Vaccination in adolescent population

The roll out for vaccinating adolescent population started last 15 October 2021 initially for those with co-morbidities and expanded to include all adolescents (with or without co-morbidities) on 02 November 2021.




Figure 3. Report distribution by age and gender

As of 05 December 2021, 1,757 reports were received. Figure 3 shows the distribution of cases by gender and age.

Out of the 1,757 reports, 66 reports were tagged as serious, 1,670 are non-serious, and 15 reports with no tag whether serious or non-serious. The most common reported reactions are dizziness, vaccination/injection site pain, blood pressure increased, headache, and pyrexia.

Pfizer and Moderna are the only vaccines with EUA for the adolescent population. Although, very rare cases of myocarditis and pericarditis have been observed in these vaccines, the benefits of using such vaccines still outweigh the risks. There are two (2) cases of myocarditis that have been reported in adolescent population. Causal link of such cases to the vaccination are being reviewed.

Booster shots

The roll out of vaccination for the booster shots (third or additional doses) started last 17 November 2021 initially for healthcare workers. This was followed by senior citizens, immunocompromised, and individuals with comorbidities at high risk of developing severe COVID-19 on 22 November 2021. In early December, general population who completed their vaccine doses may also receive booster shots.

Individuals eligible for booster shots have the option of receiving a [homologous or heterologous booster](#) dose given that they have completed their primary dose series (at least six months after getting the second dose of either CoronaVac, COVID-19 Vaccine AstraZeneca, Sputnik V, Comirnaty, or COVID-19 Vaccine Moderna; at least 3 months for Janssen COVID-19 vaccine)

As of 05 December 2021, 735 suspected adverse reaction reports were received, six (6) reports was tagged as serious while the rest are non-serious. The most common reported adverse reaction includes vaccination/injection site pain, fever, headache, blood pressure increased, chills, and malaise.

National Vaccination Day

The *Bayanihan, Bakunahan* National Vaccination Days were held last 29 November to 02 December 2021. A total of 1,043 reports were received out of more than 8 million doses administered. Nine (9) serious adverse reaction were reported while the rest are non-serious. The top reported reactions were blood pressure increased, dizziness, headache, syncope, and pyrexia.

Reports of suspected serious adverse reaction

Adverse reactions experienced after vaccination are considered serious when it resulted to any of the following criteria:

- In-patient hospitalization/prolongation of existing hospitalization

- Significant disability/incapacity
- Life-threatening (e.g. anaphylaxis) and death
- Birth defect or congenital malformations
- Considered to be medically important event

Hypersensitivity including severe allergic reactions

Severe allergic reactions have been reported on the use of COVID-19 vaccines including CoronaVac, COVID-19 Vaccine AstraZeneca, Sputnik V, Comirnaty, Janssen COVID-19 Vaccine, and COVID-19 Vaccine Sinopharm. It only occurs in a few vaccinated individuals. It usually happens in people with a history of severe vaccine reactions. Severe allergic reactions (anaphylaxis) generally occur soon after vaccination and are usually managed with Epinephrine in combination with other medicines. Thus, vaccinees are observed for at least 15 minutes after receiving their vaccine. Epinephrine is readily available in all vaccination sites in case of anaphylaxis.

The proportion of reported side effects of severe allergic reactions to COVID-19 vaccines proved to be statistically rare as the number of vaccinated populations increases. The current reporting rate for anaphylaxis is 3.19 per million doses administered.

Increased blood pressure

Blood pressure (BP) increased has been continuously reported as one of the top adverse reactions to all vaccine platforms. Monitoring BP has been part of the screening process for COVID-19 vaccination program in the country. The program recommends monitoring BP only in vaccine recipients with a history of hypertension, symptomatic hypertension, and based on the clinical judgement of the physician on the vaccination site. This is in relation to the recommendations of [the Philippine Heart Association and Philippine Society of Hypertension on elevated blood pressure readings during COVID-19 vaccination](#).

According to PRESYON 4 (Philippine Heart Association Report on the Study of Hypertension), a nationwide hypertension survey conducted in January to April 2021, the prevalence of hypertension in the Philippines alarmingly increased to 37% in 2021 among adults 18 years old and above from 28% (2013). Out of this 37%, 19% are aware of having hypertension while 18% are unaware. The BP control rate, with or without medications, is 36%. Only about 25% of hypertensive individuals monitor blood pressure at home.³ This study explains the increase in blood pressure observed in most vaccinated individuals.

³ Sison, J.A. (2021, May). Press Conference on PRESYON 4 – Nationwide 2021 Hypertension Survey Results [Video file]. Retrieved from <https://www.facebook.com/philheart.org/videos/159433679504182/>

Thrombosis-thrombocytopenia syndrome

Thrombosis-thrombocytopenia syndrome (TTS) are cases of unusual blood clots with low blood platelets. Following cases of TTS from other countries, COVID-19 Vaccine AstraZeneca and Janssen COVID-19 Vaccine revised their label to include warnings related to thrombosis with thrombocytopenia, a very rare side effect following vaccination.

Vaccinated individuals should watch out for the said adverse event and seek immediate medical assistance if they experience any sign of blood clots and low blood platelet such as:⁴

- shortness of breath
- chest pain
- leg swelling
- persistent abdominal (belly) pain
- neurological symptoms, such as severe and persistent headaches or blurred vision
- tiny blood spots under the skin beyond the site of the injection

Nine (9) cases of thrombosis have been reported. Causal link of such cases to the vaccination are being reviewed. The FDA together with the Epidemiology Bureau of the DOH shall continue to monitor vaccine safety ensuring that the benefits always outweigh the risks.

⁴ AstraZeneca's COVID-19 vaccine: EMA finds possible link to very rare cases of unusual blood clots with low blood platelets
<https://www.ema.europa.eu/en/news/astrazenecas-covid-19-vaccine-ema-finds-possible-link-very-rare-cases-unusual-blood-clots-low-blood>

Confirmed COVID-19 infections

There were 2,622 confirmed reports of COVID-19 infections. Most of the reported infections were asymptomatic cases. There were 180 severe cases which resulted to a fatal outcome. Most of the fatal reports have not yet completed their vaccination course. Upon assessment, these cases were not related to the use of the vaccine, but these were actual COVID-19 natural infections.

The vaccines currently being used in the COVID-19 vaccination program are non-replicating viral vector, inactivated, and mRNA vaccines. It does not contain any live virus and therefore does not cause COVID-19 infection in vaccine recipients.

Inflammation of the heart

Myocarditis is an inflammation of the heart muscle that may present as chest pain, palpitations, arrhythmias, and/or symptoms of heart failure while pericarditis is an inflammation of the pericardial sac that surrounds the heart and fixes it to the mediastinum. Cases of myocarditis and pericarditis on the use of mRNA vaccine, such as Comirnaty and COVID-19 Vaccine Moderna, have been reported in many countries including the US, UK, Germany, and Israel. Most of the cases are young male. The US FDA announced the revision of fact sheets for Comirnaty and COVID-19 Vaccine Moderna suggesting increased risk of myocarditis and pericarditis following vaccination. EMA's safety committee has also concluded that myocarditis and pericarditis can occur in very rare cases following Comirnaty and COVID-19 Vaccine Moderna.

Vaccinated individuals should watch out for the said adverse event and seek immediate medical assistance if they experience the following symptoms after vaccination:⁵

- breathlessness
- a forceful heartbeat that may be irregular
- chest pain

Ten (10) cases of myocarditis and one (1) case of pericarditis have been reported. Causal link of such cases to the vaccination are being reviewed. The FDA together with the Epidemiology

Bureau of the DOH shall continue to monitor vaccine safety ensuring that the benefits always outweigh the risks.

⁵ *Comirnaty and Spikevax: possible link to very rare cases of myocarditis and pericarditis*
<https://www.ema.europa.eu/en/news/comirnaty-spikevax-possible-link-very-rare-cases-myocarditis-pericarditis>

Capillary Leak Syndrome

Capillary leak syndrome is a very rare, serious condition that causes fluid leakage from small blood vessels (capillary), resulting in swelling mainly in the arms and legs, low blood pressure, thickening of the blood, and low blood levels of albumin. Several cases were reported on the use of COVID-19 Vaccine AstraZeneca and Janssen COVID-19 Vaccine. The EMA's safety committee recommended contraindication in individuals with previous capillary leak syndrome and inclusion of capillary leak syndrome as a new side effect in the product information for both products.

Vaccinated individuals should watch out for the said adverse event and seek immediate medical assistance if they experience the following symptoms days after vaccination, which may occur together with feeling faint (due to low blood pressure):⁶

- rapid swelling of the arms and legs
- sudden weight gain

No case of capillary leak syndrome has been reported on the use of COVID-19 vaccines in the Philippines as of this time. The FDA together with the Epidemiology Bureau of the DOH shall continue to monitor vaccine safety ensuring that the benefits always outweigh the risks.

⁶ *COVID-19 Vaccine Janssen: Contraindication in individuals with previous capillary leak syndrome and update on thrombosis with thrombocytopenia syndrome*
<https://www.ema.europa.eu/en/medicines/dhpc/covid-19-vaccine-janssen-contraindication-individuals-previous-capillary-leak-syndrome-update>

Guillain-Barré syndrome

Guillain-Barré syndrome (GBS) is a rare, autoimmune disorder in which a person's own immune system damages the nerves, causing muscle weakness and sometimes paralysis. An increased risk for GBS has been observed following vaccination with Janssen COVID-19 Vaccine in the US. The US FDA has announced the revision of fact sheets for Janssen COVID-19 Vaccine to include the observed risk for GBS. EMA's safety committee considered that a causal relationship between Janssen COVID-19 Vaccine and GBS is possible. COVID-19 Vaccine AstraZeneca already updated their product information.

Vaccinated individuals should watch out for the said adverse event and seek immediate medical assistance if they experience signs and symptoms suggestive of GBS such as:⁷

- double vision or difficulty moving eyes
- difficulty swallowing, speaking, or chewing
- coordination problems and unsteadiness
- difficulty walking
- tingling sensations in the hands and feet

- weakness in the limbs, chest, or face
- problems with bladder control and bowel function

Fifteen (15) cases of GBS have been reported. Four (4) cases have been assessed as product related reactions (as per published literature), three (3) cases are indeterminate meaning there is insufficient evidence that the vaccine caused the reaction, and eight (8) are currently being reviewed. The FDA together with the Epidemiology Bureau of the DOH shall continue to monitor vaccine safety ensuring that the benefits always outweigh the risks.

⁷ COVID-19 Vaccine Janssen: Guillain-Barré syndrome listed as a very rare side effect
<https://www.ema.europa.eu/en/news/covid-19-vaccine-janssen-guillain-barre-syndrome-listed-very-rare-side-effect>

Bell's palsy

Bell's palsy is a form of temporary facial paralysis or weakness on one side of the face. It results from dysfunction of facial nerve which directs the muscles on one side of the face. Cases have been reported in a number of people in Hong Kong, Canada, and UK on the use of CoronaVac, Comirnaty, and COVID-19 Vaccine Moderna. The overall number of these reports is relatively small. In relation to this, [CoronaVac vaccination fact sheet was revised to include bell's palsy as a very rare adverse reaction](#) in Hong Kong while Comirnaty product information was revised in Canada. COVID-19 Vaccine Moderna already contains this safety information.

Vaccinated individuals should watch out for the said adverse event and seek immediate medical assistance if they experience any combination of the following symptoms: ⁸

- uncoordinated movement of the muscles that control facial expressions, such as smiling, squinting, blinking, or closing the eyelid
- loss of feeling in the face
- headache
- tearing from the eye
- drooling
- lost sense of taste on the front two-thirds of the tongue
- hypersensitivity to sound in the one ear
- inability to close an eye on one side of the face

Twenty-two (22) cases of bell's palsy have been reported. Two (2) case have been assessed as product related reactions (as per published literature), four (4) cases are indeterminate meaning there is insufficient evidence that the vaccine caused the reaction, two (2) as coincidental or not related to the vaccine, and 14 are currently being reviewed. The FDA together with the Epidemiology Bureau of the DOH shall continue to monitor vaccine safety ensuring that the benefits always outweigh the risks.

⁸ Health Canada updates Pfizer-BioNTech COVID-19 vaccine label to reflect very rare reports of Bell's Palsy
<https://healthycanadians.gc.ca/recall-alert-rappel-avis/hc-sc/2021/76203a-eng.php>

Cases of hospitalization

One of the criteria for serious adverse reaction is hospitalization or extended hospital stay. Reports of adverse reaction that results in hospitalization does not necessarily mean that vaccine caused the reaction. An Expert Committee reviews and assesses whether the

vaccine caused the reaction. Based on the reports received, the hospitalization reporting rate is 2.65 per 100,000 doses administered. Commonly reported causes of hospitalization include pyrexia, cough, dyspnea, and headache.

Reports involving death

As of 05 December 2021, 1,551 fatal events were received. Reports of fatal events does not necessarily mean that the vaccine caused the events. Underlying conditions or pre-existing medical conditions causing fatal events are usually coincidental on the use of the vaccine. It is expected that reports of fatal events will rise as the vaccination program covers more people including those with undiagnosed illness, underlying comorbidities, and pre-existing medical conditions.

The vaccinees reported to have fatal events were aged 13 years and above. The mean age of the fatal cases was 62.8 years. 66.15% (1,026) of the fatal cases were from age group 60 years and above, 23.15% (359) from age group of 40-59 years, 9.16% (142) from age group 18-39 years, 0.39% (6) from age group 12-17 years of age and 1.16% (18) were not identified to what age group they are classified.

Most of these events occurred in persons with multiple existing comorbidities. These include cardiovascular diseases, ischemic heart diseases, cerebrovascular diseases, cancer, diabetes, and infections including pneumonia. There were cases of confirmed COVID-19 infections leading to severe cases with fatal outcomes.

Number of suspected adverse reactions per category

A total of 80,644 case reports consisting of 174,327 suspected adverse reactions were received from the start of the vaccination program. More than one suspected adverse reaction may be reported in a single case. Suspected adverse reactions were coded using the Medical Dictionary for Regulatory Activities (MedDRA) terminology to allow international comparison of reports.

The data presented below are categorized by System Organ Class (SOC), the highest in the hierarchy of MedDRA. They are grouped by manifestation site (e.g. gastrointestinal, cardiac) and etiology (e.g. infections, examinations).

Reactions to inactivated vaccines

- CoronaVac
- COVID-19 Vaccine Sinopharm

Classification	Number of suspected reactions
General symptoms & reactions in the administration site <i>E.g. Pain and reaction in the injection site, chills, discomfort, fever, fatigue</i>	12,773
Cardiac symptoms <i>E.g. Palpitations, bradycardia, tachycardia</i>	887

Erratum (28 February 2024): Total number of reports with fatal outcome was corrected from 1,538 to 1,551.

Ear symptoms <i>E.g. Ear swelling, vertigo, tinnitus, ear discomfort</i>	52
Endocrine symptoms <i>E.g. Adrenal insufficiency, goiter, thyroid symptoms</i>	3
Examinations <i>E.g. Increased blood pressure, increased heart rate, blood glucose increased, SARS-CoV-2 test</i>	12,101
Eye symptoms <i>E.g. Eye itchiness, blurred vision, eye pain, eye swelling</i>	265
Gastrointestinal symptoms <i>E.g. Abdominal pain, diarrhea, nausea, vomiting, dry mouth, lip swelling</i>	3,280
Hepatobiliary symptoms <i>E.g. Jaundice</i>	5
Immune system symptoms <i>E.g. Allergic reactions, hypersensitivity</i>	198
Infections <i>E.g. Cold symptoms, rhinitis</i>	3,290
Metabolism and nutrition-related symptoms <i>E.g. Decreased appetite, increased appetite, starvation, dehydration</i>	432
Musculoskeletal symptoms <i>E.g. Back pain, joint pain, pain in extremities, muscle pain, muscle spasms</i>	2,063
Neoplasm <i>E.g. Liver cancer, endometrial cancer, uterine myoma</i>	3
Neurological symptoms <i>E.g. Dizziness, headache, syncope</i>	8,492
Pregnancy, puerperium, and perinatal conditions <i>E.g. Abortion, hemorrhage</i>	3
Procedural symptoms <i>E.g. Procedural hypertension, vaccination adverse reaction</i>	84
Psychiatric symptoms <i>E.g. Feeling anxious, insomnia, nervousness, disorientation</i>	156
Renal and urinary symptoms <i>E.g. Urine coloring yellow, urine frequency</i>	45
Reproductive symptoms <i>E.g. Vaginal bleeding, vaginal spotting</i>	50
Respiratory symptoms <i>E.g. Cough, nasal congestion, throat irritation</i>	5,940
Skin symptoms <i>E.g. Cold sweat, rash, redness</i>	4,173
Social circumstances <i>E.g. Hearing disability, walking disability</i>	3
Surgical and medical procedures <i>E.g. Tumor debulking, nasolabial flap</i>	2
Symptoms in blood and lymphatic system <i>E.g. Pain in the lymph nodes</i>	37
Vascular symptoms <i>E.g. Flushes, low blood pressure</i>	395




Figure 4. Suspected adverse reaction distribution by SOC for inactivated vaccine

As shown in Figure 4, the SOC which consists of the greatest number of reports were examinations (11,977), followed by general symptoms and reactions in the administration site (9,483), neurological symptoms (7,119), respiratory symptoms (4,514), skin symptoms (3,145), infections (2,880), gastrointestinal symptoms (2,474), musculoskeletal symptoms (1,802), cardiac symptoms (858), and metabolism and nutrition-related symptoms (429).

The top reported events for CoronaVac are:

- blood pressure increased (38.26%)
- pyrexia (13.09%)
- headache (12.81%)
- vaccination/injection site pain (10.42%)
- cough (9.41%)
- rash (6.82%)
- dizziness (6.76%)
- nasopharyngitis (5.59%)
- malaise (4.95%)
- dyspnea (4.51%)

The top reported events for COVID-19 Vaccine Sinopharm are:

- pyrexia (27.92%)
- cough (14.94%), dizziness (14.94%)

- headache (14.29%)
- rash (9.74%)
- blood pressure increased (8.44%)
- nasopharyngitis (7.79%)
- dyspnea (6.49%),
- arthralgia (5.84%), chills (5.84%), fatigue (5.84%), nausea (5.84%)
- pruritus (5.19%)
- decreases appetite (4.55%), diarrhea (4.55%), hypoesthesia (4.55%)

Reactions to non-replicating viral vector vaccines

- COVID-19 vaccine AstraZeneca
- Sputnik V
- Janssen COVID-19 Vaccine

Classification	Number of suspected reactions
General symptoms & reactions in the administration site <i>E.g. Pain and reaction in the injection site, chills, discomfort, fever, fatigue</i>	44,586
Cardiac symptoms <i>E.g. Palpitations, bradycardia, tachycardia</i>	752
Congenital, familial, and genetic disorder <i>E.g. Polycystic kidney</i>	1
Ear symptoms <i>E.g. Ear swelling, vertigo, tinnitus, ear discomfort</i>	64
Endocrine symptoms <i>E.g. Adrenal insufficiency, goiter, thyroid symptoms</i>	3
Examinations <i>E.g. Increased blood pressure, increased heart rate, blood glucose increased, SARS-CoV-2 test</i>	6,925
Eye symptoms <i>E.g. Eye itchiness, blurred vision, eye pain, eye swelling</i>	493
Gastrointestinal symptoms <i>E.g. Abdominal pain, diarrhea, nausea, vomiting, lip swelling</i>	4,828
Hepatobiliary symptoms <i>E.g. Jaundice</i>	5
Immune system symptoms <i>E.g. Allergic reactions, hypersensitivity</i>	278
Infections <i>E.g. Cold symptoms, rhinitis</i>	2,872
Metabolism and nutrition-related symptoms <i>E.g. Decreased appetite, increased appetite, starvation, dehydration</i>	739
Musculoskeletal symptoms <i>E.g. Back pain, joint pain, pain in extremities, muscle pain, muscle spasms</i>	9,076
Neoplasms <i>E.g. Liver cancer, endometrial cancer, uterine myoma</i>	2
Neurological symptoms <i>E.g. Dizziness, headache, syncope</i>	16,665
Pregnancy, puerperium, and perinatal conditions <i>E.g. Abortion, hemorrhage</i>	9

Procedural symptoms <i>E.g. Procedural hypertension, vaccination adverse reaction</i>	158
Psychiatric symptoms <i>E.g. Feeling anxious, insomnia, nervousness, disorientation</i>	110
Renal and urinary symptoms <i>E.g. Urine coloring yellow, urine frequency</i>	42
Reproductive symptoms <i>E.g. Vaginal bleeding, vaginal spotting</i>	57
Respiratory symptoms <i>E.g. Cough, nasal congestion, throat irritation</i>	4,443
Skin symptoms <i>E.g. Cold sweat, rash, redness</i>	3,587
Social circumstances <i>E.g. Hearing disability, walking disability</i>	2
Surgical and medical procedures <i>E.g. Tumor debulking, nasolabial flap</i>	5
Symptoms in blood and lymphatic system <i>E.g. Pain in the lymph nodes</i>	69
Vascular symptoms <i>E.g. Flushes, low blood pressure</i>	383




Figure 5. Suspected adverse reaction distribution by SOC for viral vector vaccines

As shown in Figure 5, the SOC which consists of the greatest number of reports were general symptoms and reactions in the administration site (23,473), followed by neurological symptoms (14,505), musculoskeletal symptoms (7,353), examinations (6,881), gastrointestinal symptoms (3,711), respiratory symptoms (3,543), skin symptoms (2,767), infections (2,532), metabolism and nutrition-related symptom (732), and cardiac symptoms (720).

The top reported events for COVID-19 Vaccine AstraZeneca are:

- pyrexia (40.05%)
- headache (33.62%)
- vaccination/injection site pain (23.15%)
- malaise (21.33%)
- chills (16.20%)
- blood pressure increased (15.72%)
- myalgia (15.52%)
- fatigue (11.65%)
- arthralgia (7.68%)
- dizziness (6.44%)

The top reported events for Sputnik V are:

- blood pressure increased (55.54%)
- pyrexia (12.65%)
- headache (8.47%)
- cough (6.39%)
- rash (5.22%)
- vaccination/injection site pain (4.95%)
- nasopharyngitis (4.82%)
- dizziness (4.69%)
- heart rate increased (4.56%)
- COVID-19 (3.52%)

The top reported events for Janssen COVID-19 Vaccine are:

- blood pressure increased (27.64%)
- pyrexia (20.23%)
- vaccination/injection site pain (16.93%)
- headache (11.43%)
- cough (10.96%)
- dyspnea (6.25%)
- hypoesthesia (5.70%)
- malaise (5.62%)
- dizziness (4.79%)
- rash (4.73%)

Reactions to mRNA vaccines

- Comirnaty
- COVID-19 Vaccine Moderna

Classification	Number of suspected reactions
General symptoms & reactions in the administration site <i>E.g. Pain and reaction in the injection site, chills, discomfort, fever, fatigue</i>	7,652
Cardiac symptoms <i>E.g. Palpitations, bradycardia, tachycardia</i>	505
Ear symptoms <i>E.g. Ear swelling, vertigo, tinnitus, ear discomfort</i>	30
Endocrine symptoms <i>E.g. Adrenal insufficiency, goiter, thyroid symptoms</i>	1
Examinations <i>E.g. Increased blood pressure, increased heart rate, blood glucose increased, SARS-CoV-2 test</i>	3,612
Eye symptoms <i>E.g. Eye itchiness, blurred vision, eye pain, eye swelling</i>	155
Gastrointestinal symptoms <i>E.g. Abdominal pain, diarrhea, nausea, vomiting, lip swelling</i>	1,302
Immune system symptoms <i>E.g. Allergic reactions, hypersensitivity</i>	89
Infections <i>E.g. Cold symptoms, rhinitis</i>	803
Metabolism and nutrition-related symptoms <i>E.g. Decreased appetite, increased appetite, starvation, dehydration</i>	140
Musculoskeletal symptoms <i>E.g. Back pain, joint pain, pain in extremities, muscle pain, muscle spasms</i>	1,146
Neoplasms <i>E.g. Liver cancer, endometrial cancer, uterine myoma</i>	1
Neurological symptoms <i>E.g. Dizziness, headache, syncope</i>	3,953
Pregnancy, puerperium, and perinatal conditions <i>E.g. Abortion, hemorrhage</i>	5
Procedural symptoms <i>E.g. Procedural hypertension, vaccination adverse reaction</i>	44
Psychiatric symptoms <i>E.g. Feeling anxious, insomnia, nervousness, disorientation</i>	84
Renal and urinary symptoms <i>E.g. Urine coloring yellow, urine frequency</i>	25
Reproductive symptoms <i>E.g. Vaginal bleeding, vaginal spotting</i>	41
Respiratory symptoms <i>E.g. Cough, nasal congestion, throat irritation</i>	1,750
Skin symptoms <i>E.g. Cold sweat, rash, redness</i>	1,812
Social circumstances <i>E.g. Hearing disability, walking disability</i>	2

Surgical and medical procedure <i>E.g. Chemotherapy</i>	2
Symptoms in blood and lymphatic system <i>E.g. Pain in the lymph nodes</i>	60
Vascular symptoms <i>E.g. Flushes, low blood pressure</i>	187




Figure 6. Suspected adverse reaction distribution by SOC for mRNA vaccine

As shown in Figure 6, the SOC which consists of the greatest number of reports were general symptoms and reactions in the administration site (4,868), followed by examinations (3,563), neurological symptoms (3,297), skin symptoms (1,387), respiratory symptoms (1,378), gastrointestinal symptoms (982), musculoskeletal symptoms (947), infections (735), cardiac symptoms (496), and vascular symptoms (181).

The top reported events for Comirnaty are:

- blood pressure increased (29.54%)
- pyrexia (16.98%)
- vaccination/injection site pain (13.92%)
- headache (12.82%)
- dizziness (10.79%)
- rash (7.72%)
- cough (6.18%)
- malaise (5.13%)

- dyspnea (4.40%)
- nasopharyngitis (4.10%)

The top reported events for COVID-19 vaccine Moderna are:

- pyrexia (28.39%)
- vaccination/injection site pain (23.25%)
- blood pressure increased (22.66%)
- headache (17.92%)
- chills (8.47%)
- dizziness (7.82%)
- rash (7.27%)
- cough (7.05%)
- malaise (6.47%)
- myalgia (5.33%)

Outcome of suspected adverse reactions

The outcome of cases of suspected adverse reactions to COVID-19 vaccines is shown in Figure 7. Overall, most of the reported cases have *recovered/resolved* (76.84%), although there were few cases who have *recovered but with sequelae* (0.04%). Over 12% of the cases are *recovering/resolving* while 1% have *not recovered/not resolved* at the time of reporting. A proportion of 1.92% were reported with fatal outcomes as discussed in the section reports involving death.




Figure 7. Case outcome

Reporting of suspected adverse reactions following vaccination

Individuals who have received their COVID-19 vaccination shots can report any suspected adverse reaction to any of the following:

- Immunization site where you were vaccinated

- Directly to the vaccine manufacturer or emergency use authorization holder
 - [Sinovac – CoronaVac](#)
 - [AstraZeneca – COVID-19 Vaccine AstraZeneca](#)
 - [Gamaleya – Sputnik V](#)
 - [Pfizer – Comirnaty](#)
 - [Zuellig – COVID-19 Vaccine Moderna](#)
 - Johnson & Johnson – Janssen COVID-19 Vaccine
 - Faberco – Covovax
- [FDA online reporting system](#)

Kindly **report only to one** of the above to avoid duplication of reports.